




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G01N 33/53, C12N 15/90, 5/12, C07K 16/18, 14/51		A1	(11) International Publication Number: WO 00/62065
			(43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/09349 (22) International Filing Date: 7 April 2000 (07.04.00) (30) Priority Data: 60/128,468 9 April 1999 (09.04.99) US (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, THE NATIONAL INSTITUTES OF HEALTH [US/US]; Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FISHER, Larry, W. [US/US]; 16713 Briardale Road, Derwood, MD 20855 (US). FEDARKO, Neal, S. [US/US]; 7138 Honeyladen Place, Columbia, MD 21045 (US). YOUNG, Marian, F. [US/US]; 2112 Larier Drive, Silver Spring, MD 20910 (US). (74) Agent: NOONAN, William, D.; Klarquist, Sparkman, Campbell, Leigh & Winston, LLP, Suite 1600 - One World Trade Center, 121 SW Salmon Street, Portland, OR 97204 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: COMPLEX FORMED BY N-LINKED GLYCOPROTEINS (SIBLINGS) AND FACTOR H

(57) Abstract

The invention provides methods and compositions for exploiting the discovery that members of the small integrin-binding ligand, n-linked glycoproteins family termed SIBLINGS bind to complement Factor H, and moreover that SIBLINGS proteins, such as BSP, exist in relatively acidic forms. The methods provided can be used to detect SIBLINGS proteins in samples from subjects that are suspected of having tumors or abnormal bone turnover. The invention also provides methods of using SIBLINGS proteins to protect cells from complement mediated lysis. Finally, the discovery allows for the creation of specific binding agents that facilitate the detection of SIBLINGS proteins when they are associated with Factor H.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

FIELD OF THE INVENTION

The present invention relates to methods of detecting and utilizing small integrin-binding ligand, N-linked glycoproteins (SIBLINGS), as well as the production and use of recombinant SIBLINGS proteins that can be used as controls for such detection methods.

A. Factor H

Factor H, a 150-kDa protein, is a key regulatory braking mechanism in normal and alternate complement-mediated cell lysis. It dissociates and thereby inactivates the assembled C3 convertase, serves as an essential accelerator of Factor I-mediated cleavage of C3b to iC3b, and sterically inhibits C5 binding to C3b (a prerequisite step for terminal pathway activation). The salient structural features of Factor H include 20 short consensus repeats (SCRs) that contain four cysteine residues forming two disulfide bonds per repeat. In addition, each SCR contains at least one conserved tryptophan residue per repeat, and Factor H is known to interact with several sialic acid-containing proteins.

B. Molecular Characterization of BSP, A Member of the SIBLINGS Protein Family

Bone sialoprotein (BSP), also briefly known as BSP-II, is a phosphorylated and sulfated glycoprotein that is associated with most normal and many pathological mineralized matrices. It is a small (approximate $M_r=75,000$) integrin-binding protein that supports cell attachment *in vitro* through both RGD-dependent and RGD-independent mechanisms and has a high affinity for hydroxyapatite. BSP is a member of the family of small integrin-binding ligand, N-linked glycoproteins (SIBLINGs) family that also includes osteopontin (OPN),

dentin matrix protein (DMP1) and dentin sialophosphoprotein (DSPP). All have similar gene structures and are clustered on human chromosome 4.

Bone sialoprotein (BSP) constitutes about 10-15% of the non-collagenous proteins found in the mineralized compartment of young bone (Fisher et al., *J. Biol. Chem.* **258**:12723-727, 1983). Immunolocalization and *in situ* studies have shown BSP to be produced by osteoblasts, osteocytes, and osteoclasts (the multinucleated cells that resorb bone) (Bianco et al., *Calcif. Tissue Int.* **49**:421-426, 1991). The areas richest in BSP are the collagen-poor matrix found between areas of new bone, where bone is developing or turning over. Outside of bone, BSP has been found in other mineralized tissues, such as dentin (Fisher et al., *J. Biol. Chem.* **258**:12723-727, 1983), cementum (MacNeil et al., *J. Bone Min. Res.* **9**:1597-1606, 1994), and calcifying cartilage of the growth plate (Bianco et al., *Calcif. Tissue Int.* **49**:421-426, 1991). Trophoblasts of the developing placenta also express high levels of BSP (Bianco et al., *Calcified Tissue International*, **49**:421-426, 1991). While placental tissue is not usually considered to be a mineralized tissue, late term human placentas have hydroxyapatite crystals associated with the aging trophoblasts.

Using human BSP as a model (Fisher et al., *J. Biol. Chem.* **265**:2347-2351, 1990), the protein is first made as a 317 amino acid, 35,000 Da protein. A 16 amino acid leader peptide is removed during synthesis. BSP has no disulfide bonds and it is nearly uniformly hydrophilic along its length, indicating that the protein is likely to be an extended rod in solution. There are three regions particularly rich in glutamic acids residues ("polyglutamic acid domains") that have long been thought to govern the high affinity of this protein for hydroxyapatite. Recent work with recombinant fragments, however, shows that BSP's ability to bind strongly to apatite is found throughout its length (Stubbs et al., *Bone Miner. Res.* **12**:1210-1222, 1997).

Human BSP contains four consensus sequences for N-linked oligosaccharides, three of which are conserved for all mammalian species known to date. These N-linked and the many O-linked oligosaccharides make up approximately 50% of the mass of BSP as it is secreted into the human bone

matrix (Fisher et al., *J Biol. Chem.* **258**:12723-727, 1983). Tyrosine sulfation and serine/threonine phosphorylation make up the remainder of the known post translational modifications. There are three tyrosine-rich domains in BSP, the last two of which flank the RGD domain and are subject to sulfation. The presence or
5 absence of the sulfate groups does not appear to change the ability of fibroblasts to attach in a simple *in vitro* assay (Mintz et al., *J. Biol. Chem.* **269**:4845-4852, 1994).

The cDNA BSP sequences for rat (Oldberg et al., *J. Biol. Chem.* **263**:19430-19432, 1988), human (Fisher et al., *J. Biol. Chem.* **265**:2347-2351, 1990), mouse (Young et al., *Mamm. Genome* **5**:108-111, 1994), cow (Chenu et al., *J. Bone Miner. Res.* **9**:417-421, 1994), hamster (Sasaguri et al., Direct
10 submission to GenBank, Accession number U65889, 1996) and chicken (Yang et al., *J. Bone Miner. Res.* **10**:632-640, 1995) have been published. The human (Kerr, J.M., Fisher, L.W., Termine, J.D., Wang, M.G., McBride, O.W. and
15 Young, M.F. *Genomics* **17**:408-415, 1993) and chicken (Yang, R. and Gerstenfeld, L.C., *J. Cell. Biochem.* **64**:77-93, 1997) genes have also been published. The human IBSP gene maps very close to two other members of this family, within 340 kb of SPPI (osteopontin), and within 150 kb of DMP1 with the order being: cen-DMP1-IBSP-SPPI-tel on chromosome 4 (Aplin et al., *Genomics*
20 **30**:347-349, 1995; Crosby et al., *Genome* **7**:149-151, 1996). Mouse Ibsp is on the homologous region of chromosome 5 at 56.0 (Young et al., *Mamm. Genome* **5**:108-111, 1994). Other members of the family are also encoded on chromosome 4, for example dentin phosphoprotein and dentin sialoprotein (DSPP) are cleavage products expressed from a single transcript coded by a gene on human
25 chromosome 4 (MacDougall et al., *J Biol. Chem.* **272**(2):835, 1997).

C. Detection of Human BSP

The detection of human BSP in biological samples has been accomplished using polyclonal antibodies directed towards denatured whole BSP, non-denatured
30 whole BSP, or synthetic fragments of BSP. Certain tumors have been found to ectopically express BSP. For example, BSP has been found to be expressed by

breast cancer tissue, prostate cancer tissue, lung cancer tissue and thyroid cancer tissue (Bellahcene et al., *Cancer Research* **54**:823-826, 1994; Bellahcene et al., *Calcif. Tissue Int.* **61**:183-188, 1998; Bellahcene et al., *Calcif. Tissue Int.* **61**:183-188, 1997; and Bellahcene et al., *Thyroid* **8**:637-641, 1998 respectively).

- 5 Additional studies have shown that BSP mRNA levels are increased in human breast cancer tissue as well as cell lines derived from breast cancer tissue (Bellahcene et al., *Laboratory Investigation* **75**:203-210, 1996).

The detection of BSP in various tissue samples described above has been accomplished through the use of polyclonal antibodies directed to either whole
10 human BSP (LF-6) or synthetic fragments of human BSP, such as the synthetic fragment comprising amino acids 277-294 (LF-83).

An increased level of BSP in serum has been correlated with the presence of hyperparathyroidism, Paget's disease, multiple myeloma and breast cancer (Seibel, et al., *J. Clinical Endocrinology and Metabolism*, **81**:3289-294, 1996).

- 15 Elevated levels of serum BSP have also been detected in subjects suffering from rheumatoid arthritis (Mansson et al., *J. Clin. Invest.* **95**:1071-1077, 1995).

SUMMARY OF THE INVENTION

The present invention stems from the discovery that members of the
20 SIBLINGS family of proteins bind to Factor H and that these proteins can confer resistance to complement mediated lysis. This discovery also allows for the creation of assays that more accurately determine the total quantity of SIBLINGS proteins in samples, and for the creation of methods of utilizing SIBLINGS proteins to inhibit complement formation.

- 25 The present invention also takes advantage of the recognition that the population of BSP (one member of the SIBLINGS family of proteins) in serum contains two distinct sub-populations. The first sub-population of BSP is relatively non-acidic, and it is the predominant sub-population found in normal subjects. The second sub-population of BSP is relatively acidic and it is found
30 predominantly in subjects with various types of tumors. Accordingly, the invention also provides methods of detecting the relatively acidic BSP and the

relatively non-acidic BSP. Furthermore, these methods can be practiced for example, by using monoclonal antibodies specific for either the relatively acidic BSP or by simply separating the different forms of BSP based upon their ionic charges.

5 One embodiment of the invention involves detecting at least one member of the SIBLINGS family of proteins in a manner that Factor H does not inhibit the detection of the SIBLINGS protein. The detection of the SIBLINGS protein can occur *in vivo* or *in vitro*, for example in the context of a sample.

In another embodiment the invention provides methods of detecting the
10 SIBLINGS protein by separating the SIBLINGS protein from the SIBLINGS/Factor H complex and then detecting the SIBLINGS protein. Separation of the SIBLINGS protein from Factor H can be accomplished in a variety of ways, for example by heating the sample in the presence of reducing agents such as DTT (1,4-dithiothreitol) and β -mercaptoethanol. These methods
15 can also incorporate the use of denaturants such as urea, SDS, and formamide.

Factor H can also be separated from the SIBLINGS protein based upon the differences between the ionic charge of the SIBLINGS protein and Factor H. Factor H can also be separated from the SIBLINGS protein by reacting a SIBLINGS/Factor H complex with a specific binding agent.

20 The specific binding agent can be specific for the portion of the SIBLINGS protein that is exposed in the SIBLINGS/Factor H complex or it can be specific for the SIBLINGS/Factor H complex itself. When the specific binding agent is specific for the SIBLINGS/Factor H complex itself it is not necessary to separate the SIBLINGS protein from Factor H prior to detection. Furthermore, in some
25 instances the specific binding agent can serve to separate and identify the SIBLINGS protein simultaneously.

The invention also provides methods of making specific binding agents that detect the exposed portion of SIBLINGS proteins when they are complexed to Factor H as well as specific binding agents that specifically detect the
30 SIBLINGS/Factor H complex. Accordingly, the invention also provides compositions containing these specific binding agents.

The methods provided by the invention are particularly useful for screening for the presence or the recurrence of a tumor, such as a liquid tumor, prostate tumor, thyroid tumor, lung tumor, or breast tumor that is associated with abnormal SIBLINGS protein production. The methods can be practiced for example, by
5 testing samples of body fluids, such as blood, serum, or saliva. In some instances the tumor may produce a relatively acidic SIBLINGS protein, such as BSP, and therefore, the invention provides methods of detecting the relatively acidic BSP population.

Accordingly, the invention provides transformed organisms that expresses
10 recombinant SIBLINGS proteins such as BSP, OPN, DSPP, and DMP1 (rBSP, rOPN, rDSPP, and rDMP1, respectively). This organism can express full length SIBLINGS proteins or fragments thereof, more specifically the organism can express the BSP molecule encoded by the construct shown in SEQ ID NO: 8. This recombinantly produced BSP can then be purified for use in a variety of different
15 applications.

In another embodiment the invention provides a method of conferring protection from a complement mediated immune response a subject. This method involves providing a reservoir or other supply in the subjects body from which a SIBLINGS protein can be dispersed to interfere with complement mediated lysis
20 and inflammation. For example a cell or an implant can be contacted (for example coated) with a SIBLINGS protein. When such a coated implant is introduced into the body, it will provoke less of an inflammatory response. In the case of a cell, a recombinant nucleic acid sequence can be introduced via transformation and the cell will then express the SIBLINGS protein. This embodiment is useful for
25 protecting cells that are grafted onto foreign tissue or for protecting bone marrow cells that are being introduced into a foreign host. Alternatively, a reservoir of SIBLINGS protein is placed in the body for metered dispensing of the protein in therapeutic dosages.

In another embodiment of the invention the SIBLINGS protein can be
30 detected in a subject that is suspected of having abnormal bone turnover, for example abnormal bone turnover associated with osteoporosis.

These and other aspects of the invention will become readily apparent in light of the description provided below.

BRIEF DESCRIPTION OF THE FIGURES

5 **Figs. 1A-C** are graphs which show the binding of BSP to Factor H as detected with HP-conjugated avidin after separation on a molecular sieve column. Fig. 1A shows a chromatograph of biotinylated BSP. Fig. 1B shows biotinylated BSP preincubated with normal human serum, and Fig. 1C shows biotinylated BSP preincubated with Factor H.

10 **Figs. 2A-C** are graphs which show the separation of BSP from Factor H after treatment with formamide, DTT, and heat, as detected with LF-100 antisera after separation on a molecular sieve column. Figure 2A shows a chromatograph of normal human serum (NHS). Figure 2B shows biotinylated BSP-(btBSP) incubated with normal human serum and Fig. 2C shows normal human serum after
15 incubation with DTT in the presence of formamide and heat.

Figs. 3A and 3B are graphs which show the result of a titration of Factor H with BSP. Fig. 3A shows relative fluorescence of Factor H, and Factor H complexed with BSP. Fig. 3B shows the fractional acceptor saturation (f_a) vs. the molar ratio of BSP to Factor H (C_s), the latter of which was constant throughout
20 the assay. This figure shows that BSP binds 1:1 with Factor H and has a binding constant of ≤ 1 nM. Similar results were seen with OPN and DMP1.

Figs. 4A-F are graphs which show profiles of serum samples generated by separating the two sub-populations of BSP on a Toyopearl QAE column. Fig. 4A and B show two normal sera, Fig. 4C shows the profile from a subject known to
25 have a thyroid tumor. Fig. 4D shows the profile from a subject known to have a lung tumor. Fig. 4E shows the profile from a subject known have a breast tumor, and Fig. 4F shows the profile from a subject known to have a prostate tumor.

Figs. 5A-E: show profiles from 5 different subjects that were known to have breast cancer.

Fig. 6 shows the level of serum BSP detected in samples taken from 32 individuals under the age of 50, and the serum BSP levels from 36 individuals over the age of 50.

Fig. 7 shows the level of serum BSP detected in samples taken from normal subjects, subjects known to have prostate cancer, and subjects known to have breast cancer.

Figs. 8A and 8B are graphs which show that recombinantly produced BSP and OPN protect MEL cells from complement mediated lysis. MEL cells were rinsed three times with GVB-MgEGTA buffer, resuspended in GVB-MgEGTA at a density of 5×10^6 cells/mL, and incubated at 37°C with different concentrations of normal human serum. After 2 hrs, cells were harvested for trypan blue exclusion assay. The thiazolium blue assay was carried out at identical serum dilutions and cell viability was determined by absorbance at 560 nm. Each data point represents the average of three measurements. The error bars represent the standard deviation of the mean (A). MEL cells in GVB-MgEGTA buffer were incubated with 10 µg/mL of either rBSP or rOPN for 10 minutes at 37°C. Normal human serum was then added at a dilution of 1:10 and the cells were returned to 37°C, incubated for 2 hours and cell viability was determined by trypan blue and thiazolyl blue (MTT) reduction assays (B). Data from 3-7 separate experiments, with treatment replicates in triplicate, was combined to yield mean values. Error bars represent the standard error of the mean. N = number of experiments combined.

Fig. 9 is a bar graph which shows the involvement of integrin and CD44 in protection from complement mediated lysis. MEL cells prepared as in Fig. 8 were treated with normal rBSP (10 µg/mL), rBSP whose RGD sequence had been mutated to KAE (10 µg/mL), and either GRGDS (SEQ ID NO: 10) peptide (400 µM) followed by rBSP or an αVβ3 antibody (1:4000) followed by rBSP for 10 minutes prior to the addition of normal human serum. The cells were then incubated for 10 minutes after which cell viability determined using the MTT assay (A). A cohort of MEL cells were pretreated with rOPN (10 µg/mL) alone, GRGDS (SEQ ID NO: 10) peptide followed by rOPN, the αVβ3 antibody

followed by rOPN, or an anti-CD44 antibody (Chemicon, Co.) followed by rOPN, or hyaluronan (HA) followed by rOPN (B). Cells were then treated with normal human serum and viability assayed as in Fig. 8. Treatment of MEL cells with a pre-formed complex of either BSP-Factor H ([BSP + fH]) or OPN-Factor H ([OPN + fH]) abolished the protection from 23 complement-mediated lysis. The data represents the mean and standard error of the mean for three separate experiments. Statistical significance was determined by analysis of variance. Percent cell viability was determined using A560 absorbance values of various conditions and a control where no serum had been added (100 % viable). The cross-hatched region represents that range of values observed when normal human serum (1:10) alone was added (maximal cell death).

Figs 10A and 10B are bar graphs which show that human cancer cell lines can be protected from complement mediated lysis through the addition of SIBLINGS proteins. Human cancer cell lines, MCF-7 (A) and U-266 (B) cells were rinsed three time with GVB-MgEGTA buffer and subsequently treated exactly as in the MEL cell complement-mediated cell lysis assay described above except complement active guinea pig serum (GPS) was substituted for human serum. Three different dilutions of GPS were assayed. For cultures pretreated with 10 $\mu\text{g/mL}$ BSP or OPN, the dilution of GPS used was 1:5. Cell viability was determined by both trypan blue exclusion (solid bar) and MTT (open bar) assay. The bars represent the mean values of triplicate samples and the error bars are the standard deviation values.

Figs 11A and 11B are bar graphs which show the separation of DMP1 from Factor H on a molecular sieve column after treatment with formamide, DTT, and heat, as detected by DMP1 specific polyclonal antisera. Fig. 11A shows normal human serum (NHS) and Fig. 11B shows NHS that has been treated.

SEQUENCE LISTINGS

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids. Only one strand of each nucleic acid sequence is

shown, but the complementary strand is understood as included by any reference to the displayed strand.

SEQ ID NO: 1: shows the amino acid sequence of human BSP.

SEQ ID NOS: 2-7: show various amino acid sequences which have been
5 used as immunogens for the creation of BSP specific antibodies.

SEQ ID NO: 8: shows a cDNA sequence encoding human BSP.

SEQ ID NO: 9: shows an amino acid sequence that was used as an immunogen for the creation of BSP specific antibodies.

SEQ ID NO: 10: shows the amino acid sequence of a peptide that was
10 used to compete with proteins from the SIBLINGS family of proteins for integrin type binding.

SEQ ID NO: 11: shows the nucleic acid sequence of an OPN encoding sequence (Accession No. J04765).

SEQ ID NO: 12: shows the amino acid sequence derived from the
15 sequence shown SEQ ID NO:11.

SEQ ID NO: 13: shows the nucleic acid sequence of an OPN encoding sequence (Accession No. X13694).

SEQ ID NO: 14: shows the amino acid sequence derived from the sequence shown SEQ ID NO:13.

SEQ ID NO: 15: shows the nucleic acid sequence of a DMP1 encoding
20 sequence (Accession No. NM_004407).

SEQ ID NO: 16: shows the amino acid sequence derived from the sequence shown in SEQ ID NO: 15.

SEQ ID NO: 17: shows the nucleic acid sequence of a DSPP encoding
25 sequence (Accession No. AF163151).

SEQ ID NO: 18: shows the amino acid sequence derived from the sequence shown in SEQ ID NO: 17.

DETAILED DESCRIPTION

I. Abbreviations and Definitions

Abnormal: Deviation from normal characteristics. Normal characteristics can be found in a control, a standard for a population, etc. For instance, where
5 the abnormal condition is a disease condition, such as a tumor, a few appropriate sources of normal characteristics might include an individual who is not suffering from the disease (*e.g.*, breast cancer), a population standard of individuals believed not to be suffering from the disease, etc.

Likewise, abnormal may refer to a condition that is associated with a
10 disease. The term "associated with" includes an increased risk of developing the disease as well as the disease itself. For instance, a certain abnormality (such as an increase in the expression of one or more SIBLINGS proteins) can be described as being associated with the biological condition of breast cancer; thus, the abnormality is predictive both of an increased risk of developing breast cancer and
15 of the presence of breast cancer.

Abnormal protein expression, such as abnormal SIBLINGS protein expression, refers to expression of a protein that is in some manner different from expression of the protein in a normal (wildtype) situation. This includes but is not necessarily limited to: (1) a mutation in the protein such that one or more of the
20 amino acid residues is different; (2) a short deletion or addition of one or a few amino acid residues to the sequence of the protein; (3) a longer deletion or addition of amino acid residues, such that an entire protein domain or sub-domain is removed or added; (4) expression of an increased amount of the protein, compared to a control or standard amount; (5) expression of an decreased amount of the
25 protein, compared to a control or standard amount; (6) alteration of the subcellular localization or targeting of the protein; (7) alteration of the temporally regulated expression of the protein (such that the protein is expressed when it normally would not be, or alternatively is not expressed when it normally would be); (8) alteration in post translational processing; and (9) alteration of the localized (*e.g.*,
30 organ or tissue specific) expression of the protein (such that the protein is not

expressed where it would normally be expressed or is expressed where it normally would not be expressed), each compared to a control or standard.

Controls or standards appropriate for comparison to a sample, for the determination of abnormality, include samples believed to be normal as well as
5 laboratory values, even though possibly arbitrarily set, keeping in mind that such values may vary from laboratory to laboratory. Laboratory standards and values may be set based on a known or determined population value and may be supplied in the format of a graph or table that permits easy comparison of measured, experimentally determined values.

10 **Implant:** A non-self object that is placed in contact with a subject. For example, the object can be an artificial limb, an intraosseous, or intravascular implant or a dental prosthesis. Such objects can comprise molecules that are generated through biosynthetic pathways, such as hydroxyapatite, or collagen, or
15 the object can contain molecules that are generated through non-biosynthetic means, such as metal alloys or plastics. Furthermore, such objects can contain a mixture of molecules such that some of the molecules are biosynthetically generated and some of the molecules are generated through non-biosynthetic means.

Small Integrin-Binding Ligand, N-Linked Glycoproteins (SIBLINGS):

20 The family of secreted phosphoproteins that includes, for example, bone sialoprotein (BSP), osteopontin (OSP), dentin matrix protein (DMP1), and dentin sialophosphoprotein (DSPP). It is also likely that other member of the SIBLINGS family of proteins will be identified in the future, and that these additional members of the SIBLINGS family will also bind to Factor H. Hence, when the
25 term SIBLINGS protein is used it refers to not only to the four family members described above, but also to additional family members that will be identified in the future.

The genes encoding the four members of the SIBLINGS family are located on human chromosome 4. Members of the SIBLINGS family of protein are
30 shown, *infra*, to be capable of binding Factor H. The SIBLINGS/Factor H complex can cause the SIBLINGS protein to be nearly undetectable unless the

complex is first disrupted. Hence, the discovery that members of the SIBLINGS family of proteins binds to Factor H allows for more accurate and sensitive quantification of the concentration of SIBLINGS proteins. The discovery that SIBLINGS proteins bind Factor H has also led to the development of methods of using SIBLINGS proteins to selectively protect cells from complement mediated lysis.

Biologically Active SIBLINGS Proteins: Biologically active SIBLINGS proteins are characterized by their ability to protect cells from complement mediated lysis. This activity can be readily assessed using the MEL cell assay described below. These proteins can be "derived" from SIBLINGS proteins such that the endogenous SIBLINGS protein's amino acid sequence is altered to contain amino acid substitutions, deletions and/or additions. If, however, the endogenous SIBLINGS protein sequence is altered it will continue to display its ability to inhibit complement-mediated lysis and the derived SIBLINGS protein sequence will be additionally characterized by having at least 50%, 60%, 70%, 80%, or 90% sequence identity to the endogenous sequence.

As mentioned above, the inhibition of complement-mediated lysis can be detected using the MEL cell based assay described below. A SIBLINGS protein will be found to be "biologically active" if when tested in the MEL assay it confers protection to cells compared to a control sample that does not contain the SIBLINGS protein.

BSP: The normal endogenous population of bone sialoprotein. This population contains a wide range of BSP proteins, some that are not extensively post-translationally modified, as well as some of which are extensively post-translationally modified. Furthermore, this normal population contains within it at least two distinct sub-populations. The first sub-population is relatively non-acidic, and it is the sub-population that is most prominent in the serum of normal subjects (subjects without tumors). The second sub-population of BSP is relatively acidic and found in subjects with tumors, for example in subjects with breast, lung, prostate, and thyroid tumors.

Relatively Acidic BSP: The term relatively acidic BSP refers to a sub-population of BSP found primarily in the serum of subjects who have various types of tumors. There are multiple methods that can be used to detect the presence of this sub-population in a subject. These methods include column chromatography methods and electrophoresis. Therefore, for clarity the relatively acidic BSP is defined through its behavior on the specific ion exchange column assay described in section IV, below. Briefly, this section details the use of a Toyopearl QAE column for the separation of the relatively non-acidic BSP from the relatively acidic BSP. Characterized in this way, and using the assay provided, the relatively acidic BSP will elute from the column at a salt concentration of greater than 0.5 M NaCl.

While not wishing to be bound to a particular theory, it is believed that the relatively acidic BSP is post-translationally modified to contain sialic acids, phosphate, and/or sulfate. In that case, the relative concentration of each sub-population of phosphorylated or sulfated BSP can be determined by assays that quantify the concentration of sulfur and/or phosphate.

Specific binding agent: An agent that binds substantially only to a defined target. Thus, a SIBLINGS protein specific binding agent binds substantially only a particular SIBLINGS protein. For example, the term "BSP specific binding agent" includes anti-BSP and other agents that bind substantially only to BSP. Similarly, the term "relatively acidic BSP specific binding agent" includes antibodies that are specific for the relatively acidic BSP and other agents that bind substantially only to the relatively acidic BSP. The term "relatively non-acidic BSP specific binding agent" includes antibodies that are specific for the relatively acidic BSP and other agents that bind substantially only to the relatively non-acidic BSP. The term "exposed portion of SIBLINGS protein specific binding agent" includes antibodies and other agents that are specific for the exposed portion of a SIBLINGS protein when it is complexed with Factor H.

The discussion below specifically mentions anti-BSP antibodies, however it also applies to anti-relatively acidic BSP antibodies, anti-relatively non-acidic BSP antibodies, as well as anti-exposed portion of SIBLINGS protein antibodies.

The term anti-SIBLINGS protein antibodies encompasses monoclonal and polyclonal antibodies that are specific for SIBLINGS proteins, i.e. which bind substantially only to BSP when assessed using the methods described below, as well as immunologically effective portions (fragments) thereof. Preferably, the anti-SIBLINGS protein antibodies used in the present invention are monoclonal antibodies (or immunologically effective portions thereof) and may also be humanized monoclonal antibodies (or immunologically effective portions thereof). Immunologically effective portions of monoclonal antibodies include Fab, Fab', F(ab')₂, Fabc and Fv portions (for a review, see Better and Horowitz, *Methods Enzymol.* **178**:476-496, 1989). Anti-BSP antibodies may also be produced using standard procedures described in a number of texts, including Antibodies, A Laboratory Manual by Harlow and Lane, Cold Spring Harbor Laboratory (1988).

The determination that a particular agent binds substantially only to the desired immunogen may readily be made by using or adapting routine procedures. One suitable *in vitro* assay makes use of the Western blotting procedure (described in many standard texts, including Antibodies, A Laboratory Manual by Harlow and Lane, Cold Spring Harbor Laboratory, 1988). Western blotting may be used to determine that a given specific binding agent, such as an anti-SIBLINGS protein monoclonal antibody, binds substantially only to a SIBLINGS protein, as described in Fisher et al., *JBC*, **262**:9702-08, 1987.

Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed host cell. A vector may include nucleic acid sequences that permit it to replicate in the host cell, such as an origin of replication. A vector may also include one or more selectable marker genes and other genetic elements known in the art.

Transformed: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Sample: The term sample includes any artificially generated or biological specimen that is of interest and is being tested for the presence of SIBLINGS proteins. More specifically, the term biological specimen includes any specimen that was generated by a living cell. Therefore, samples derived from cells in a tissue culture setting, including the cells themselves fall within the definition or biological specimen. Furthermore, samples taken from a subject, such as tissue samples and bodily fluids (such a blood plasma or serum) also fall within the definition of biological specimen.

Tumor: Tumors are abnormal growths which can be either malignant or benign, solid or liquid (for example, hematogenous). This term particularly includes malignant tumors which can be either solid (such as a breast or prostate carcinoma) or non-solid (such as a leukemia). Tumors can also be further divided into subtypes, such as adenocarcinomas (e.g. of the breast, prostate or lung).

Metastasis: A tumor implant discontinuous with a primary tumor. Any cancer may metastasize to bone, but metastases from carcinomas are the most common. There is a subset of carcinomas that is clinically recognized as more likely to metastasize to bone, and this subset includes carcinomas of the breast, lung, prostate, kidney and thyroid. Even if a tumor is not presently recognized as being particularly likely to metastasize to bone, it can be so categorized by a diagnosis in a particular subject that a skeletal metastasis has occurred (for example by scintigram and biopsy), or by studies of populations of subjects presenting with a particular histological sub-type of cancer.

Isolated: An isolated biological component (such as a nucleic acid or protein) has been substantially separated or purified away from other biological components in the biological specimen in which the component naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids and proteins which have been isolated thus include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids and proteins.

Purified: The term purified does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified BSP protein preparation is one in which the BSP is more enriched than the protein is in its natural environment. That environment being either in serum or within a cell.

- 5 Preferably, a preparation of BSP is purified such that BSP represents at least 50% of the total protein content of the preparation.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a
10 promoter is operably linked to a coding sequence, if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in the same reading frame.

Recombinant: A recombinant nucleic acid is one that has a sequence that
15 is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques.

20 **Subject:** This term includes both human and non-human mammals. Similarly, the term subject includes both human and veterinary subjects.

Sequence identity: The term "sequence identity" is used to describe the similarity between two nucleic acid sequences or between two amino acid sequences. Sequence identity is typically expressed in terms of percentage
25 identity; the higher the percentage, the more similar the two sequences.

Methods for aligning sequences for comparison purposes are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444-2448, 1988; Higgins and Sharp, *Gene* 73:237-244, 1988; Higgins and Sharp, *CABIOS* 5:151-153, 1989; Corpet et al., *Nucleic Acids Research* 16:10881-10890, 1988;
30

Huang et al., *Computer Applications in the Biosciences* 8:155-165, 1992; and Pearson et al., *Methods in Molecular Biology* 24:307-331, 1994; Altschul et al., *J. Mol. Biol.* 215:403-410, 1990, presents a detailed consideration of sequence alignment methods and homology calculations.

5 The NCBI Basic Local Alignment Search Tool (BLAST™, Altschul et al., *J. Mol. Biol.* 215:403-410, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the Internet, for use in connection with the sequence-analysis programs blastp, blastn, blastx, tblastn and tblastx.

10 For comparisons of amino acid sequences of greater than about 30 amino acids, the "Blast 2 sequences" function in the BLAST™ program is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per-residue gap cost of 1). When aligning short peptides (fewer than about 30 amino acids), the alignment should be performed using the Blast 2
15 sequences function, employing the PAM30 matrix set to default parameters (open gap 9, extension gap 1 penalties). Proteins having even greater similarity to the reference sequences will show increasing percentage identities when assessed by this method, such as at least 45%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% sequence identity.

20 A first nucleic acid is "substantially similar" to a second nucleic acid if, when optimally aligned (with appropriate nucleotide insertions or deletions) with the other nucleic acid (or its complementary strand), nucleotide sequence identity occurs in at least about 60%, 75%, 80%, 85%, 90% or 95% of the nucleotide bases. (As used herein, "optimally aligned" sequences exhibit a maximal possible
25 sequence identity). Sequence similarity can be determined by comparing the nucleotide sequences of two nucleic acids using the BLAST™ sequence analysis software (blastn) available from The National Center for Biotechnology Information. Such comparisons may be made using the software set to default settings (expect = 10, filter = default, descriptions = 500 pairwise,
30 alignments = 500, alignment view = standard, gap existence cost = 11, per residue existence = 1, per residue gap cost = 0.85). Similarly, a first polypeptide

is substantially similar to a second polypeptide if it shows sequence identity of at least about 75%-90% or greater when optimally aligned and compared using BLASTTM software (blastp) using default settings.

Conservative amino acid substitutions: Conservative amino acid substitutions usually have minimal impact on the activity of the resultant protein. Such substitutions are described below.

Conservative substitutions replace one amino acid with another amino acid that is similar in size, hydrophobicity, etc. Such substitutions generally are conservative when it is desired to finely modulate the characteristics of the protein. Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include: Ser for Ala; Lys for Arg; Gln or His for Asn; Glu for Asp; Ser for Cys; Asn for Gln; Asp for Glu; Pro for Gly; Asn or Gln for His; Leu or Val for Ile; Ile or Val for Leu; Arg or Gln for Lys; Leu or Ile for Met; Met, Leu or Tyr for Phe; Thr for Ser; Ser for Thr; Tyr for Trp; Trp or Phe for Tyr; and Ile or Leu for Val.

More substantial changes in enzymatic function or other features may be obtained by selecting substitutions that are less conservative than those above, i.e., selecting residues that differ more significantly in their effect on maintaining: (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Substitutions that generally produce the greatest changes in protein properties are those in which: (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl, or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine. The effects of these amino acid substitutions or deletions or additions may be assessed for SIBLINGS protein derivatives by analyzing the

ability of the respective modified polypeptide to inhibit complement-mediated cell lysis.

Variant SIBLINGS protein encoding cDNAs or genes may be produced by standard DNA mutagenesis techniques, for example, M13 primer mutagenesis.

- 5 Details of these techniques are provided in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Ch. 15, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al. (ed.) *Current Protocols in Molecular Biology*, Greene Publishing and Wiley-Interscience, New York (with periodic updates), 1987. By the use of such
- 10 techniques, variants may be created that differ slightly from the endogenous SIBLINGS protein encoding cDNA or gene sequences, yet that still encode a biologically active SIBLINGS protein. DNA molecules and nucleotide sequences that are derivatives of those specifically disclosed herein and that differ from those disclosed by the deletion, addition, or substitution of nucleotides while still
- 15 encoding a biologically active SIBLINGS protein are comprehended by this invention. In their simplest form, such variants may differ from the disclosed sequences by alteration of the coding region to fit the codon usage bias of the particular organism into which the molecule is to be introduced.

- Additionally, as mentioned above new members of the SIBLINGS family of
- 20 proteins can be identified by searching various DNA databases, such as the database maintained at the National Center for Biotechnology Information in Bethesda, MD. Searches can be done such that sequences that are conserved between existing members of the SIBLINGS family of proteins are used to identify other members of the family. Once identified these members can be used to the
- 25 assays described below to determine if they maintain SIBLINGS protein biological activity.

Throughout the specification and claims, reference to the singular (such as "a" or "the") includes the plural, unless clearly indicated otherwise by context.

II. Identification and Characterization of SIBLINGS/Factor H Complex

Initial experimentation, which is described below, led to the identification of a BSP/Factor H complex. The population of BSP in serum from subjects displaying tumors was additionally found to contain a sub-population of relatively acidic BSP. Subsequent experimentation showed that both OPN and DMP1 bound to Factor H and it is likely that the detection of these SIBLINGS proteins, as well as other members of the protein family, will be enhanced by disrupting the SIBLINGS/Factor H complex prior to quantifying protein levels.

A. Identification and Characterization of the BSP/Factor H Complex

BSP is an acidic protein, containing not only many acidic amino acids (particularly glutamic acid) but also phosphate, sulfated tyrosines and oligosaccharides, as well as many sialic acid groups. As such it should bind to an anion exchange column under conditions of low salt (<0.1 M NaCl) and neutral pH (6-8). This is indeed the case, and BSP from bone extracts and from culture medium has been found to elute off of a DEAE column at about 0.4 M NaCl in 7 M urea at pH 6.

Surprisingly, serum BSP failed to behave in the same manner as BSP from bone extracts and cell culture medium as described above. In fact, injecting normal human serum onto a DEAE or QAE HPLC column resulted in all of the detectable BSP eluting in the unbound fraction, therefore it behaved as a neutral or positively charged protein. This indicated that the BSP was bound to something that masked its many negative charges, hence attempts were made to disrupt the complex. These attempts involved using denaturants, such as 50% formamide or 8 M urea, reducing agents such as 1 mM DTT or 10 mM β -mercaptoethanol, and heating to 100°C for 5 minutes. None of these approaches was independently successful in disrupting the complex, as indicated by the failure of the BSP to bind to an anion exchange column.

However, a combination of a denaturant, a reducing agent and heating finally did separate the BSP from the complex. This separation was noted by the increased binding of the BSP to an anion exchange column with subsequent elution

in a salt gradient and detection with polyclonal antisera. The anti-sera used was raised in rabbit against human BSP or fragments thereof, and termed LF-6, LF-83, LF-84, LF-100, LF-101, LF-119, LF-120, and LF-125 (see Table 2), which were tested for specificity via either ELISA assay or Western blotting. (Acta Orthop. Scand, (Suppl. 266) 66:61-65, 1995. The LF-142 (Table 2) polyclonal antisera was raised in rabbit against a synthetic BSP peptide which contained several sulfated tyrosines.

The previously uncharacterized protein described above, that was found to mask the otherwise acidic BSP, was subsequently identified as being Factor H. The identity of the protein was determined by several different assay techniques, and these techniques, as well as the results are described below.

1) Factor H specific monoclonal antibodies were obtained from Quidel, San Diego, California, USA. These antibodies were used to precipitate Factor H from human sera. One sample was left untreated while another was heated to 100°C for 5 minutes in the presence of 1 mM DTT and 50% formamide. These samples were then separated on a SDS gel and subjected to western blotting analysis using a BSP specific antibody LF-100. Results from the Western blot analysis showed a high molecular weight band ($M_r \sim 200,000$ Da) in the lane without DTT, and the more typical BSP size of 75,000 Da in the lane treated with DTT and formamide.

Furthermore, a similar high molecular weight band was also seen when a Western blot was performed using as a sample purified human Factor H (Quidel) complexed with BSP. This artificially created complex was immunoprecipitated by the same Factor H antisera (see above), and analyzed with the same results as with the whole human serum described above. This showed that BSP was bound to Factor H.

2) Biotinylated human BSP was added to normal human serum and the material was chromatographed on a calibrated molecular sieve column (Superose 6, Pharmacia, Fig. 1B). A separate chromatogram with the biotinylated BSP alone was also generated. The biotinylated products were subsequently detected using HRP-conjugated avidin on a 96-well direct ELISA (Fig. 1A). The BSP alone chromatographed to the expected approximate 75,000 Da position, while the BSP

added to the human serum chromatographed to a position consistent with being complexed to Factor H (about 200,000 Da, Fig. 1B, detected with HRP-conjugated avidin). Furthermore, no 75,000 Da, (i.e. free) BSP could be detected, suggesting that all was bound to the Factor H (Factor H is found at concentration of about 500 µg/mL human serum). An additional chromatograph was generated with biotinylated BSP added to Factor H (Fig. 1C). This chromatograph showed that the product eluted at the same size at the sample containing biotinylated BSP and normal human serum (Fig. 1B).

The effect of treatment with DTT and 50% formamide was observed using a molecular sieve column and detecting the products with the polyclonal antibody LF-100. This antibody was previously found to bind to both the bound and unbound form of BSP, however, this binding is probably due to the fact that the SIBLINGS/Factor H complex is at least partially disrupted. Results showed that the BSP from untreated normal human serum eluted at about 200,000 Da (Fig. 2A). Similarly, biotinylated BSP with normal human serum showed a major peak at about 200,000 Da, the expected size of the BSP/Factor H complex, and a second small peak about 75,000 Da (Fig. 2B). Finally, a sample of normal human serum incubated in the presence of 50% formamide and 1 mM DTT at 100°C for five minutes resulted in a peak corresponding to BSP (Fig. 2C).

3) Biotinylated human BSP bound normally to the anion exchange column. Addition of human serum prior to loading on the anion exchange column caused the biotin activity to elute in the unbound fraction. Pretreating with purified human Factor H gave the same results. These results showed that Factor H could render the BSP unable to bind to the column, a property of the BSP found naturally in human serum.

4) Fluorescence spectroscopic analysis of the tryptophanes of Factor H (BSP does not contain tryptophane) during the binding of BSP shows that the two rapidly form a strong interaction that is stoichiometric (1:1) and saturable (Fig. 3A and 3B).

B. Characterization of OPN and DMP1/Factor H Binding

The salient structural features of Factor H include 20 short consensus repeats (SCRs) that contain four cysteine residues forming two disulfide bonds per repeat. In addition, each SCR contains one conserved tryptophan residue per repeat and Factor H is known to interact with several sialic acid-containing proteins. Therefore, binding interactions between members of the SIBLINGS family of proteins and Factor H can be studied by steady state fluorescence.

Steady state fluorescence studies were accomplished by titrating purified human complement Factor H with rBSP, rOPN, and rDMP1 followed by excitation at 295 nm and monitoring emission between 300 and 450 nm. The emission profile of Factor H alone yields a peak at 347 nm. The addition of rBSP, rOPN, or rDMP1 in nm increments causes a relative fluorescent intensity quenching. Conversion of the fluorescent intensity titration into a binding curve by determining the fraction of binding sites occupied as the fractional change in fluorescence quenching at 347 nm yields a saturable binding curve. By steady state fluorescence, the binding of BSP, OPN, and DMP1, by Factor H are saturable and possess a 1:1 stoichiometry and the binding constants are in the nM range. Given that normal serum concentration of Factor H is ~ 0.5 mg/mL it is probable that virtually all BSP, OPN, and DMP1 in serum will be complexed with Factor H.

III. Detection of Elevated Levels of Members of the SIBLINGS Family of Proteins in Serum

Recent observations have shown that BSP and OPN are expressed by malignant tissue. BSP is expressed in primary breast cancers (Gillespie et al., *Int. J. Cancer* **73**(6):813-815, 1997; Bellahcene et al., *Cancer Res.* **54**(11):2823-2826, 1994; Bellahcene et al., *Int. J. Cancer* **69**(4):350-353, 1996; Seibel et al., *J. Clin. Endocrinol Metab.* **81**(9):3289-3294, 1996), prostate cancer (Waltregny et al., *J. Natl. Cancer Inst.* **90**(13):1000-1008, 1998), lung cancer (Bellahcene et al., *Calcif. Tissue Int.* **61**(3):183-188, 1997), thyroid cancer (Bellahcene et al., *Thyroid* **8**(8):637-641, 1998), malignant bone disease (Chen et al., *Histochem.* **30**(1):1-6,

1998), and by neoplastic odontoblasts (van der Pluijm et al., *Cancer Res.* 56(8):1948-1955, 1996). In addition, BSP peptides are potent inhibitors of breast cancer cell adhesion to bone (van der Pluijm et al., *Lab Invest.* 77(6):665-675, 1997; Tunio et al., *Arch. Pathol. Lab. Med.* 122(12):1087-1090, 1998). OPN is
5 expressed in breast cancer (Gillespie et al., *Int. J. Cancer.* 73(6):812-815, 1997; Sung et al., *Exp. Cell Res.* 241(2):273-284, 1998; Bellahcene and Castronovo, *Bull. Cancer.* 84(1):17-24, 1997; Tuck et al., *Int. J. Cancer* 79(5):502-508, 1998), as well as in prostate cancer (Koeneman et al., *Prostate* 39(4):246-261, 1999), thyroid cancer (Wright et al., *Int. J. Cancer.* 46(1):39-49, 1990), skin
10 cancer (Senger et al., *CBiochim. Biophys. Acta* 996(1-2):43-48, 1989) and several other types of cancer (Senger et al., *Anticancer Res.* 56(8):1948-1955, 1996; Chambers et al., *Anticancer Res.* 12(1):43-47, 1992; Mevoracj et al., *J. Exp. Med.* 188(12):2313-2320, 2331, 1998). Given that these two SIBLINGS proteins are expressed in cancer tissue it is likely that other members of the family will also
15 be found to be over expressed in tumors. Hence, it is desirable to develop sensitive assays for detecting the presence of SIBLINGS proteins.

As described above, it has been shown the BSP, OPN, and DMP1 bind to Factor H. Hence, it is likely that these members of the SIBLINGS family of proteins as, well as the yet untested DPP1 protein, are bound to Factor H *in vivo*.
20 Therefore, in order to accurately detect the concentration of these proteins in serum the SIBLINGS/Factor H complex should be disrupted prior to quantifying the SIBLINGS protein level.

Serum levels of BSP have been quantified previously via radioimmunoassays (RIAs). These assays, which involve incubation for a 24 hour
25 period in the presence of antibody, have led to the determination that the circulating levels of BSP in normal subjects are between 5 ng/mL and 24 ng/mL (Karmatschek, *Clinical Chemistry* 43:2076-82, 1997). However, using the experimental procedures detailed below, which exploit the finding that BSP is at least partially undetectable when bound to Factor H, it has been determined that
30 the circulating concentration of BSP in normal subjects may extend across a much broader range, for example between 200 ng/mL serum and 1000 ng/mL serum.

Thus, the discovery that BSP is bound to Factor H allows BSP levels to be more accurately determined.

The direct detection of BSP as well as other SIBLINGS proteins in serum is hampered by the fact that it is effectively bound by Factor H and is not expected to be easily detected by most antisera. For example, serum samples where eluted from an ion exchange column, and ELISA assays (described below) were preformed using polyclonal antisera LF-6 and LF-83 (Table 2). These antisera were the same polyclonal antisera that were used by Bellahcene et al., *Cancer Research* 54:823-826, 1994; Bellahcene et al., *Calcif. Tissue Int.* 61:183-188, 1997; Bellahcene et al., *Calcif. Tissue Int.* 61:183-188, 1998a; and Bellahcene et al., *Thyroid* 8:637-641. 1998b, to detect the extopic expression of BSP in various tumor types. However, these antisera, as well as several others failed to detect the BSP/Factor H complex found in serum. Therefore, disrupting the BSP/Factor H complex prior to quantitating serum BSP levels allows for a more accurate assessment of serum BSP concentration.

There are several different methods which could be used to disrupt the SIBLINGS/Factor H complex and detect serum SIBLINGS protein levels. However, the methods used ideally take into account: 1) the properties of the SIBLINGS/Factor H complex compared to the SIBLINGS protein once it is freed from Factor H; 2) the possible problems resulting from denaturing other serum proteins, such as albumin (for example, denatured serum in 50% formamide is acceptable, but if the denaturant present is subsequently removed, the serum forms a gel even if diluted 1:10 in normal saline buffer); and 3) the SIBLINGS protein should also end up in a medium that is suitable for conducting immunoassays after the SIBLINGS/Factor H complex is disrupted.

One particularly useful method of detecting SIBLINGS proteins involves separating the SIBLINGS protein from Factor H, inactivating the contaminants, such as DTT and BME, and then diluting the sample such that the sample is in a media that allows for immunoassay techniques to be used. This assay method is more feasible, in part, because of the considerably higher concentration of SIBLINGS protein that is available after separation form

Factor H.

Another example of a method for quantifying serum levels of BSP as well as other SIBLINGS proteins involves: 1) separating the SIBLINGS/Factor H complex from most other serum components, particularly albumin; 2) disrupting the complex (by substantial separation of Factor H from the SIBLINGS protein); and 3) introducing the SIBLINGS protein into buffer conditions compatible with binding to antibodies. The initial separation step involved diluting 50 μ l serum with 150 μ l PBS. The resulting diluted serum was then placed on a disposable anion exchange column (Toyopearl QAE column available from Toso-Haas, a part of Rhom&Hass, Montgomeryville, Pennsylvania). The albumin present in the serum was then bound to the column and the Factor H- SIBLINGS protein complex passed through. The complex was then separated via incubating at 100°C in the presence of DTT and formamide. The denaturants and reducing agents were then removed by placing the sample on a sephadex G25 column available from Amersham Pharmacia, Piscataway, New Jersey, which was previously equilibrated with normal saline or some other ELISA- or RIA-compatible buffer. The eluted SIBLINGS protein was then detected by ELISA assay.

Used in this way the invention allows for the development of screening assays for the detection of SIBLINGS protein in serum. These assays involve separating the SIBLINGS protein from the SIBLINGS protein/Factor H complex and identifying the free SIBLINGS protein. Therefore, these assays can be used in conjunction with several already existing monoclonal and polyclonal antisera, as well as other antibodies developed in the future. This will allow for the more accurate assessment of SIBLINGS protein levels as well as the ability to assess the presence of several disease states in subjects. For example, the presence of tumors, Paget's disease, rheumatoid arthritis and hyperparathyroidism can be detected via the above-described assay.

A. Identification Of A Highly Acidic Form Of BSP In Subjects Known To Have Tumors

As described above, subjects with certain types of tumors have a population of BSP which ranges from a peak which elutes at a salt concentration that is

similar to that of normal subjects, to a peak which elutes later than BSP from a normal subjects. Therefore, in subjects with tumors (particularly certain types of tumors, such as those that often metastasize to bone) the serum BSP tends to be not only increased in total amount but also differs from that of a normal subject. This difference, between the BSP populations of normal subjects and subjects with tumors, can be exploited for the development of an assay for detecting tumors. One such assay is described below.

Differences in elution profiles between subjects with tumors and subjects without tumors was illustrated by taking 50 μ l of serum from normal volunteers and from patients with tumors prior to surgery (no special treatment is required for drawing or holding the serum). Subsequently, 50 μ l of fresh formamide and 1 μ l of 100 mM DTT were added to the serum sample, and it was heated in a sand-filled heating block (100°C) for 5 minutes, thereby separating the BSP from Factor H. The sample was then microfuged for 1 minute at 15,000 x g and injected onto a 1 mL packed volume Toyopearl QAE (Toso-Haas) column, that had been pre-equilibrated with 0.05 M sodium phosphate, 50% fresh formamide, pH 7.4. The column was then washed using a flow rate of 2 mL/minutes for 5 minutes in the same low salt buffer that was used for equilibration, and then the salt concentration was linearly increased to 2 M NaCl (in the same buffer, formamide is optional) over 80 minutes (Figure 4A-F). Finally, the salt concentration was at 2 M NaCl for the final 15 minutes.

Similar results were seen in five different subjects with breast tumors. The method used is described above with the exception that a steeper salt gradient was used (25 minutes) followed by 2 M NaCl for the final 15 minutes (Fig. 5A-E).

Then 100 μ l of each fraction was placed in a corresponding well of a microtiter plate. It was found that Greiner High-binding plates or Nunc Maxisorp worked well for binding both the relatively non-acidic form of BSP and the relatively acidic form of BSP. The microtiter plates were incubated for between 1 and 18 hours at 4°C. The plates were then washed three times (5 minutes each) with >200 μ L/well Tris-buffered saline plus 0.05% Tween 20 (TBS-T) and exposed to 100 μ L of rabbit polyclonal antiserum (LF-100) raised against human

BSP in TBS-T for 1 hour at room temperature. The plates were then washed again as described above, and exposed to 100 μ L 1:2000 HRP-conjugated goat anti-rabbit IgG (human serum adsorbed, Kirkegaard & Perry, Gaithersburg, Maryland) in TBS-T for one hour at room temperature. Finally, the plates were washed a
5 third time and the color was developed using 3,3', 5,5' tetramethylbenzidine and H_2O_2 for 10-30 minutes at room temperature. Color development was stopped by that addition of 25 μ L 1 N H_2SO_4 and analyzed at 450 nm.

Results from the above described HPLC separation technique indicated that the relatively acidic BSP was detected only in sera from subjects with various
10 tumors and was eluted from the column generally at salt concentrations greater than 1 M NaCl. It was readily apparent that subjects with these various tumors not only had increased levels of BSP, but also that they had a different population of BSP compared to that of normal subjects.

The use of the semi-quantitative method described above allowed for the
15 detection of a relatively non-acidic form of BSP that eluted early in the gradient (≤ 1 M NaCl). Of course, altering the chromatography procedure used, such as by changing the type of resin, could lead to elution of the peak at a different salt concentration, however, regardless of the separation method used there will be a sub-population of detected BSP which is less negatively charged, and will elute
20 separately from the more negatively charged fraction. In normal subjects this relatively less negatively charged sub-population would be the only or at least the dominant population which is detected. However, in the case of normal subjects it is still possible that there will be at least a trace amount of the relatively acidic BSP. It is believed that the relatively non-acidic BSP can be used to measure bone
25 turnover generally.

Furthermore, the relatively non-acidic BSP population has been detected in both normal subjects, as well as in subjects known to have tumors. Subjects who have prostate tumors may have elevated levels of the relatively non-acidic BSP compared to normal, which possibly indicates that there is an elevated level of
30 bone turnover.

The above-described assay is designed to detect any tumor that makes BSP and has a sufficiently high tumor load. Using the techniques described herein, it can be determined whether any particular tumor type is associated with increased BSP levels, and particularly acidic forms of BSP. Furthermore, the identification
5 of the relatively acidic form also provides a more sensitive assay than those based on merely assessing total serum BSP levels.

Furthermore, the above-described assay could be simplified so that only three steps are needed. These steps are, removal of the albumin by binding de-complexed BSP in low salt (≤ 0.2 M NaCl), eluting the relatively non-acidic BSP
10 with 1 M NaCl and subsequently eluting the relatively acidic BSP with 2 M NaCl. Samples can then be placed in a solution that allows for antibody binding and the BSP recovered can be detected by ELISA.

**B. Creation of Specific Binding Agents Which Recognize Free BSP,
As Well As Bound BSP**

15 Nine rabbit (polyclonal) antisera against human BSP or fragments (LF-6, LF-83, LF-84, LF-100, LF-101, LF-119, LF-120 and LF-125) (including peptides) have been made. These include eight polyclonal antibodies whose activity is summarized in the Acta Orthop. Scand, (Suppl. 266) 66:61-65, 1995, as well as one made since (LF-142) that has not been published. The immunogens
20 used to produce these antisera is provided below in Table 2.

TABLE 2

Gene Product	Antiserum	Antigen	Known Species
Human BSP	LF-6	human bone BSP	H,M,D,R,Mou
Human BSP	LF-83	YESENGEPRGDNYRAY ED- (LPH) (SEQ ID NO: 2)	H,M,D,R,Mou,C
Human BSP	LF-84	YESENGEPRGDNYRAY ED- (LPH) (SEQ ID NO: 2)	H,M,D,R,Mou,C
Human BSP ,	LF-100	AIQLPKKAGDIC-(LPH) (SEQ ID NO: 3)	H,M,D,R
Human BSP	LF-101	AIQLPKKAGDIC-(CSA) (SEQ ID NO: 3)	H,M,D,R,P,B
Human BSP	LF-II9	recomb.RGDdomain (AA257-317) (SEQ ID NO: 4)	H,M,D,C
Human BSP	LF-120	recomb. Frag I (AA 129--281) (SEQ ID NO: 5)	H,M,D,P,R
Human BSP	LF-125	recomb. (36-61) (SEQ ID NO: 6)	H,M,P,R,S
Human BSP	LF-142	CTVEY*EGEY*EY*TGA NDY*DNGY*EIY*ESEN Y*(SEQ ID NO: 7)	H

H=human, M=monkey, D=dog R=rat, Mou=mouse, C=chicken, P=pig,
S=sheep, B=bovine, LPH=horseshoe crab hemocyanin, CSA=chicken serum
albumin

5

*indicates sulfated tyrocine

Of these 9 polyclonal antisera only LF-100 can immunoprecipitate natural BSP, and even LF-100 only binds to a portion of the BSP (<5%) when it is complexed with Factor H in human serum as detected by Western blot analysis. The other
5 antisera precipitate little or no BSP/Factor H complex.

A similar finding was made when an artificial complex was made by incubating an excess of purified Factor H with biotinylated BSP. Once again, only the LF-100 antisera was capable of precipitating even a small portion of the BSP/Factor H complex.

10 The ability of polyclonal antisera against LF-100 to precipitate both bound and unbound BSP makes it highly likely that a monoclonal antibody to the same fragment would behave in the same manner. This monoclonal equivalent to LF-100 can be made by injecting into mice the identical or similar peptide-carrier conjugate, as discussed in Example 2. Clones derived from these mice can also be
15 screened for the ability to recognize both bound and unbound BSP.

It is also believed that a monoclonal antibody against the amino-terminus or carboxy-terminus of BSP will work because these domains may remain free of the BSP-Factor H complex.

Alternatively, a monoclonal antibody may be made by first obtaining the
20 BSP/Factor H complex, and then using the complex itself as an immunogen. The BSP/Factor H complex is obtained either through isolating it from serum or through artificially creating it using isolated Factor H and recombinant BSP. For example, 5 mg of purified human Factor H (from Quidel, LaJolla CA) can be complexed *in vitro* with 2 mg of purified human BSP from the adenoviral system
25 described below. The complex can then be passed through a QAE column without denaturants to remove any unbound BSP. The complex can then be chromatographed on a molecular sieve column (Superose 6, Pharmacia) to separate the complex from any unbound Factor H. The resulting complex is injected into mice for monoclonal production. All clones that can bind to recombinant BSP in
30 the initial screening can then be assayed to see if they can immunoprecipitate the BSP/Factor H complex. Even those that can not immunoprecipitate the intact

complex will be useful as BSP monoclonals for the detection of BSP after it has been separated from Factor H, or for use in screening for ectopic expression as described below.

The method of using the BSP/Factor H complex as an immunogen described above can also be used with other SIBLINGS protein. Thus, monoclonal antibodies which bind to other SIBLINGS/Factor H complexes can be generated.

C. Creation Of Specific Binding Agents Which Recognize The Relatively Acidic Form Of BSP

Monoclonal antibodies that recognize the relatively acidic form of BSP can be made by using the relatively acidic BSP isolated from human serum; using relatively acidic BSP isolated from tumor tissue; using synthetic peptides which are highly phosphorylated or sulfated; or by using recombinant BSP. A particularly promising method of producing monoclonal antibodies to the relatively acidic form of BSP is by complexing relatively acidic BSP to Factor H and using the complex as an immunogen. This offers the possibility of recovering an antibody which recognizes both the bound and unbound forms of the relatively acidic BSP.

Methods of obtaining the recombinant BSP for use as an immunogen are described below, and methods of obtaining the BSP/Factor H complex for use as an immunogen are described above.

The relatively acidic form of BSP can be isolated directly from human tumors using the denaturing procedures provided in Fisher et al., *J Biol. Chem.* **258**:12723-727, 1983. Furthermore, methods of generating such monoclonal antibodies are well known to those of skill in the art and these methods are additionally detailed below. Briefly, subsequent to injection of the immunogen and fusion, the clones are screened for binding to the relatively acidic form of BSP, and then subsequently screened for binding to the relatively non-acidic form of BSP. Monoclonal antibodies that specifically detect the relatively acidic form of BSP are selected for subsequent identification of this substance in biological specimens.

D. Detection of Elevated Levels of Serum BSP in Subjects known to have Breast Cancer or Prostate cancer

Sera were obtained from clinically defined normal, prostate and breast cancer patients (East Coast Biologics, Inc., North Burwick, Maine and Johns
5 Hopkins Bayview Medical Center, General Clinical Research Center) under informed consent with IRB approval and quenched samples were taken for analysis.

The day before serum was collected a 20 µg/mL solution of recombinant BSP (rBSP) was diluted out from a 1 mg/mL stock solution in 4 M guanidine HCl.
10 100 µl of the dilute BSP was added each well of a Greiner high binding (product number 655061) 96 well plate. The plate was then covered and incubated overnight at 4° C, however, shorter incubation periods are possible.

A solution of TBS-Tween containing 0.05 M Tris-HCl, pH 7.5, 0.15 M NaCl, 0.05 % Tween 20, was prepared. A blocking buffer containing Tris
15 buffered saline (TBS)+ 1% BSA + 5% powdered milk was also prepared.

Samples were diluted 1:10 in a 30% formamide 40 mM phosphate buffer, pH 7.4. The samples were then reduced with 2 mM dithiothreitol (1/100 of stock, =2.5 µL) at 100°C for 5 minutes. Residual reducing agent was inactivated by the addition of equimolar H₂O₂. BSP standards containing 2000, 400, 80, 16, 3.2,
20 0.64, and 0.13 ng/mL rBSP were then diluted out.

LF-100 antibody was then diluted 1:2,000 in TBS-Tween and 100 µl /well of Ab solution was added to polypropylene 96 well 0.5 mL volume plate (USA Scientific which is same as Nunc version). 100 µl of standard, or sample was then added to each well. The lid was then placed on the plate and reacted at room
25 temp, 1 hour.

During incubation of the samples the rBSP coated plate, described above, was removed from the cold and 250 µl blocking buffer was added to each well. The blocking buffer was allowed to react for 5 minutes and then it was removed. The plate was then washed three times with TBS-Tween.

30 100 µl reaction mixture was added to each well of the rBSP-coated plate. The plate was then incubated for 1 hour at room temp. After incubation the plate

was washed three times using a buffer containing ~250 μ L/well TBS-Tween, and then blotted and drained.

The secondary antibody (Goat anti-rabbit IgG (H&L), peroxidase-labeled antibody conjugate, that was human serum adsorbed) which is available from
5 Kirkegaard & Perry, Gaithersburg, MD, was diluted 1:20,000 in TBS-Tween. 100 μ L of the secondary antibody was then added to each well and the plates were incubated for 1 hour at room temp. After incubation, the plates were washed three times with ~250 μ L/well TBS-Tween, blotted and then drained.

The results were visualized by providing 100 μ L/well TMB (3,3', 5,5"-
10 tetramethylbenzidine) microwell peroxidase substrate (BioFX Laboratories, Randallstown, MD) and incubating the plates for 30 minutes. The reaction was then stopped through the addition of 100 μ L stop solution/well and the plates were read at 450 nM.

The results are shown in Fig. 7, and Table 3 below. The mean
15 concentration of BSP for the 32 subjects with breast cancer was 1170 ng/mL, the mean concentration of BSP for the 22 subjects tested with prostate cancer was 1320 ng/mL, and finally the mean concentration of BSP for the 72 normal subjects was 490 ng/mL. Clearly, most subjects tested in both the prostate group and the breast cancer group have elevated levels of BSP. Some have BSP levels that are
20 within the normal range, however, this is expected because not all tumors are thought to express BSP.

E. Detection of Elevated Levels of Serum BSP in Subjects over Age 50

A group of 36 individuals over the age of 50, and a group of 32 individuals
25 under the age of 50 were tested to determine their serum BSP levels. The results are shown in Fig. 6, and Table 3, below. The mean of the group under the age of 50 was clearly less than that of the over 50 group. This assay, therefore, shows that serum levels of BSP and possibly other members of the SIBLINGS family of proteins can be used to measure bone turnover or other age-related changes.

Table 3

	< 50 years Normal	> 50 years Normals	prostate	breast	All Normals
BSP mean (ng/mL)	344.266	619.787	1320	1170	490.13
Standard Deviation	41.318	179.953	289.582	364.396	192.096
Number of Individuals Sampled	32	36	32	22	68

F. Creation Of Cell Cultures Expressing Human BSP In The Highly Acidic Form

5 Full-length human BSP cDNA (GenBank accession number J05213; Seq. ID NO: 8) was excised from the original plasmid with *EcoRI* and subcloned into the *EcoRI* site of the pAC-EF1 adenovirus construct plasmid. This construct contained the EF-1 promoter and the bovine growth hormone (BGH) 3'

10 polyadenylation message stabilization sequences, which is a modification of the original pAC vector (Gluzman et al., Eucaryotic Viral Vectors, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 187-192, 1982). Selection for double cross-over within the replication-deficient adenovirus genome results in replication-deficient adenovirus that can infect most cell types but can be

15 amplified only in specific cell culture lines. Infected cells produce the human BSP, modify it according to their specific ability, and secrete the product into the culture media.

A similar cloning scheme allowed for the production of adenovirus constructs that encode OPN and DMP1 allowing choice of all lines.

20 One method of producing the relatively acidic BSP involves human marrow stromal fibroblasts. These cells in primary culture modify the BSP and can produce the BSP under serum-free conditions for several days. A small percentage of the BSP made in these marrow fibroblasts elutes in the high salt portion of the anion exchange column. Additionally, this relatively acidic BSP produced by the

25 fibroblasts infected with the adenoviral vector described above stains a darker blue

(characteristic of being more acidic) than other fractions of BSP that are eluted off of the anion exchange columns earlier, when using StainsAll dye for detection ((1-ethyl-2-[3-(1-ethylnaptho[1,2d]-thiazolin-2-ylidene)-2-methylpropenyl]naptho[1,2d]thiazolium bromide, made by Eastman Kodak, Rochester, New York).

Additional adenoviral constructs were also made that contain coding regions for OPN and DMP1. These constructs also can be used to express OPN and DMP1 in various cell types. It is also believed that SIBLINGS proteins, such as OPN and DMP1, also will be found to be postrationally modified such as has been seen in the case of BSP. Hence, the adenoviral vectors that are used to produce OPN and DMP1 can be used to produce different varieties of postrationally modified OPN and DMP1 by simply infecting different cell types. The expressed OPN and DMP1 can then be used in various assays for controls and to generate antibodies.

Transformation of normal fibroblasts with the BSP encoding vector (~ 100,000,000 cells on a total of eight 15 cm plates) allowed for the purification 20-40 mg of human BSP. Amino terminal sequencing verified that the amino-terminus corresponds to that of human BSP. Additionally, LF-83 bound to the product, hence the carboxy-terminus of the recombinant BSP is complete at least through amino acid 295 (of 317).

Furthermore, it is believed that other primary cell lines, as well as other cancer cell lines, transformed with the vector described above produce one or both sub-populations of BSP in cell culture. Several cancer cell lines are readily available from ATCC and other sources (for example: MCF-7, CRL-1435, CRL-1470, SAOS-2, MG-63 etc.).

G. Uses Of Specific Binding Agents Directed Against The Relatively Acidic BSP

BSP specific binding agents, as well as other SIBLINGS protein specific binding agents, such as those made by the methods described herein, can be used for immunostaining of biopsies and sections of various tumor types, which can assist in diagnosing and determining the severity of disease. BSP levels may also

be used to determine possible treatments that would be most suitable for a particular patient. For example, a very strong BSP positive breast cancer biopsy can be used as an indicator that more aggressive treatment is required following surgery (such as adjuvant chemotherapy or radiation therapy). Furthermore,
5 monoclonal antibodies to human BSP may also be useful as immunostaining agents for determining that an otherwise inconclusive biopsy is, in fact, cancer.

Examples of BSP specific binding agents are the 13 monoclonal antibodies that were produced through the use of a synthetic 27-amino acid peptide (256T-282N). Of the 13 monoclonal antibodies produced 4 are IgG type and 9 are IgM
10 type. This peptide was derived from the human BSP sequence, and it was designed such that it contained five tyrosine sulfates and was conjugated to a carrier protein through an amino terminal Cys (keyhole limpet hemocyanin, KLH). Monoclonal antibodies from mice injected with this construct were selected if they bound to 1) the same peptide conjugated to a different carrier protein (bovine
15 serum albumin); 2) the same peptide, but without the tyrosines sulfation conjugated to the second carrier protein; and 3) recombinant BSP made from human marrow fibroblasts infected with the virus construct described above.

H. Use Of Specific Binding Agents That Bind To The Portion Of BSP That Does Not Contribute To Increased Activity

20 This method allows for the detection of the relatively acid BSP. The BSP/Factor H complex is disrupted and the decomplexed BSP is allowed to bind to an anti-BSP antibody coated plate or bead that specifically binds to regions of BSP that are not involved in increased acidity. (Examples of such regions are epitopes with the domains used for LF-100 or LF-125, but many other likely areas can be
25 easily chosen.) Then a second binding agent can be added, such as an antiserum, lectin, or other chemical reagent (including those commercially available) which can detect various chemical groups that may contribute to the increased relative acidity of the BSP in cancer patient serum (phosphoserine, phosphothreonine, sulfated tyrosine, various sialic acids, various sulfated sugar groups, etc.). This
30 second binding agent binds to the BSP that was previously bound by the first antiserum. The second antiserum or chemical reagent is then quantified using

standard direct or indirect methods. This approach allows development of a quantitative assay more specific for the relatively acidic BSP without necessarily relying on a single antibody system that is specific to the BSP protein. For example, if the increased acidity is determined to be caused by a sialic acid group
5 the BSP can be trapped by an antiserum, such as LF-100 or a monoclonal antibody devised to that domain, and then the sialic acid can be detected and quantified. The post-translational modification itself may be common in many proteins in the serum (or urine etc.), however the detection of the modification on the captured BSP makes the assay useful for the detection of a disease.

10 IV. Inhibition of Complement Cascade via Expression of SIBLINGS

Proteins

A. Results

The key role of Factor H is the regulation of complement and alternate complement activity. Factor H can also be 'hijacked' by invading organisms to
15 permit pathogen evasion of complement mediated lysis. Pathogens such as *Streptococcus pyogenes* (Kotarsky et al., *J. Immunol.* 160(7):3349-3354, 1998; Jarvis and Vedros, *Infect. Immun.* 55(1):174-180, 1987), *Neisseria gonorrhoeae* (Ram et al., *J. Exp. Med.* 188(4):671-680, 1999; Ram et al., *J. Exp. Med.*, 187(5):743-752, 1998; Diaz and Sim, *J. Immun.* 80(3):473-482, 1995), and
20 *Echinococcus granulosus* (Diaz and Sim, *J. Immun.* 158(8):3779-3786, 1997; Wiles et al., *J. Mol. Biol.* 272(2):253-265, 1997) bind Factor H to their cell surface and are resistant to complement-mediated cell lysis. In addition, molecular mimicry of Factor H occurs when a pathogen makes a protein that is similar in sequence to Factor H to defend against attack by the host complement system as
25 described in Vaccinia virus (Isaacs et al., *Proc. Natl. Acad. Sci.* 89(2):628-632, 1992; Huemer et al., *Immunology* 79(4):639-647, 1993), Herpes simplex virus (Donelson et al., *Molecular Biol.* 91(1):51-66, 1998), and Trypanosoma cruzi (Hall and Joiner, *J. Eukaryot Microbiol.* 40(2):207-213, 1993; Kipnes and daSilva, *Braz. J. Med. Biol. Res.* 22(1):1-16, 1989; Ryden et al., *Eur. J. Biochem.*
30 184(2):331-336, 1989). The following data demonstrates that BSP, DMP1, and OPN play a similar role in tumor cell complement evasion.

The first model system employed was a murine erythroleukemia (MEL) cell line which when incubated with normal human serum can be readily assayed for alternate complement pathway (ACP)-mediated cell lysis (Varnet et al., *Curr. Opin. Cell Biol.*, 8(5):724-730, 1996). Cell survival was measured by both trypan blue dye exclusion and by thiazolyl blue (MTT) reduction by living mitochondria. Titration with dilutions of normal human serum and time courses were carried out to define optimal incubation conditions. At 1:10 dilution, human serum totally lysed the MEL cells as measured by both assay systems (Fig. 8A). The addition of purified recombinant BSP to MEL cells followed by treatment with normal human serum completely protected the cells from lysis (Fig. 8B). Treatment of MEL cells with recombinant OPN (Fig. 8B) or DMP1 also conferred protection from ACP-mediated cell lysis. The protection of MEL cells from alternate complement pathway-mediated cell lysis by BSP, OPN, and DMP1 exhibited dose related responses.

The mechanism of protection from complement-mediated lysis was investigated. Preincubation of MEL cells with rBSP in which the RGD sequence had been mutated to KAE completely removed the protective affects of this protein (Fig. 9). Furthermore, preincubation of MEL cells with either GRGDS (SEQ ID NO: 10) peptide or an $\alpha V\beta 3$ antibody (which blocks that integrin's binding activity) negated the protective effect of rBSP. Up-regulation of the $\alpha V\beta 3$ integrin has also been shown to correlate with invasive potential (Culty et al., *J. Cell Biol.* 111(6 Pt 1):2765-2774, 1990). For OPN, pretreatment with GRGDS (SEQ ID NO: 10) peptide or the $\alpha V\beta 3$ antibody reduced cell survival, though the magnitude of reduction was not as great as that seen for BSP (Fig. 9B). OPN has multiple potential receptors including $\alpha V\beta 1, \beta 3$, $\beta 5$ -containing integrin receptors as well as CD44, a receptor implicated in attachment, homing and aggregation of lymphocytes as well as neoplastic cells. Pretreatment of MEL cells with hyaluronan, a natural ligand for CD44 (Faulk and Temple, *Nature* 262(5571):799-802, 1976), as well as with an anti-CD44 antibody also reduced the protective effect of added rOPN (Fig. 9). Treatment of MEL cells with a pre-formed complex of either rBSP-Factor H or rOPN-Factor H abolished the protection from

complement-mediated lysis. For BSP, this is consistent with immunoprecipitation data that indicates that the RGD moiety is inaccessible in the solution phase Factor H complex. Hence, both of these proteins are believed to be entirely masked by Factor H shortly after being secreted by a cell.

5 Additional assays were conducted to determine if DMP1 could also confer protection against complement. DMP1 was found to protect the cells from complement-mediated lysis. Preincubation of the cells with antisera to either $\alpha V\beta 3$ integrin or CD44 diminished the DMP1 protection and together they abolished it ($p > 0.001$). Addition of the synthetic GRGDS (SEQ ID NO: 10)
10 peptide diminished but did not abolish the protective properties of DMP1. DMP1 appears to protect the cells from attack by complement by bridging the major complement control protein, Factor H, to cell surface integrins through its RGD motif and through CD44.

To verify that the protective effect of BSP and OPN is exhibited in human
15 cancer cells, MCF-7 breast cancer cells selected for non-adherent growth and U-266 myeloma cells were used in the alternate complement pathway cell lysis assay with guinea pig serum as the source of complement activity. Both cell types exhibited enhanced survival and protection from complement-mediated cell lysis when rBSP or rOPN were present (Fig 10). Increasing concentration of guinea pig
20 serum led to decreasing cell viability, while the pretreatment with either 10 $\mu\text{g/mL}$ rBSP or rOPN resulted in increased cell viability.

Collectively the above described results show that DMP1, OPN, and BSP bridge Factor H to integrins and inactivate the destructive complement pathway. These results also confirm that OPN and DMP1 bind to CD44. Furthermore, it is
25 likely that other members of the SIBLINGS family will have the same biological activity.

B. Experimental Procedures

1. Reagents

Rabbit Anti-human BSP antibodies LF-83, LF-100, LF-119, LF-120, LF-
30 125 have been previously described (Fisher et al., *Bacta Orthop. Scand. Suppl.* 266(1):61-65, 1995). Rabbit Anti-BSP peptide-derived antibody LF-142 and

mouse monoclonal antibodies LFmAb-11, LFmAb-2, LFmAb-12, and LFmAb-13 were raised against the sequence EY*EY*TGVNEY*DNGY*EIY*ESENGETP (amino acids 258 – 285; SEQ ID NO: 9) conjugated to horseshoe crab hemocyanin, where the Y* denotes tyrosine-sulfates. Normal human serum, purified human complement Factor H protein and mouse monoclonal antibody against Factor H were obtained from Quidel Corporation (San Diego, CA). Polyclonal antibodies against CD-44 and a "functional" antibody against $\alpha V\beta 3$ were obtained from Chemicon Co. (Temecula, CA). Synthetic purified glycine-arginine-aspartate-serine peptide (GRGDS; SEQ ID NO: 10) was obtained from Calbiochem-NovaBiochem Corp. (La Jolla, CA). Pre-immune serum, human serum adsorbed goat anti-rabbit IgG conjugated to horse radish peroxidase (HRP) as well as goat anti-mouse IgG conjugated to HRP were obtained from Kirkegaard & Perry (Gaithersburg, MD). HRP-conjugated streptavidin and sulfosuccinimidobiotin were obtained from Pierce Chemical Co. (Chicago, IL). Alpha-Minimal Essential Medium (MEM), Dulbecco's Modified Essential Medium (DMEM), RPMI 1640, Eagle's Minimal Essential Medium (EMEM), Earle's Balanced salt solution, Hank's balanced salt solution and heat inactivated fetal bovine serum (FBS) were obtained from BioFluids, Inc. (Rockville, MD).

2. Western Blotting

Samples diluted in gel sample buffer were resolved by SDS-PAGE 4 - 20 % gradient gels (Novex Corp, San Diego, CA), transferred to nitrocellulose following standard conditions (Towbin et al., *Proc. Natl. Acad. Sci.* **76**(9):4350-4354, 1979). Nitrocellulose membranes were rinsed with Tris-buffered saline (0.05 M Tris-HCl, pH 7.5, 0.15 M NaCl) containing 0.05 % Tween 20 (TBS-Tween). After a 1 hour incubation in blocking solution (TBS-Tween + 5 % non-fat powdered milk) at room temperature on rotary shaker, 1:5000 primary antibody was added and incubated overnight at 4°C. The nitrocellulose sheet was washed in TBS-Tween four times for 5 minutes each time with TBS-Tween and then 1:50,000 HRP conjugated second antibody in TBS-Tween + 5 % milk was added and incubated for 2 hours at room temperature. Following removal of the second antibody solution the membrane was washed three times with TBS-Tween

and rinsed a final time in enzyme substrate buffer for 5 min. Enhanced chemiluminescence reagents were employed for signal detection (Pierce Chemical Co., Chicago, IL) with x-ray film.

3. High Performance Liquid Chromatography

5 A Shimadzu LC10AS binary gradient system was employed for chromatographic separations. A 1.0 mL packed volume ToyoPearl QAE (TosoHaas, Montgomeryville, PA) column was pre-equilibrated with 0.05 M sodium phosphate, pH 7.4, containing 50 % fresh formamide. A linear salt gradient increasing to 2.0 M NaCl at 2.0 mL/minute flow rate over 50 minutes
10 was employed collecting 1 minute fractions. Size exclusion chromatography utilized a 1.0 x 30 cm Superose 6 column (Amersham Pharmacia, Piscataway, New Jersey) equilibrated in 0.05 M sodium phosphate, pH 7.4, containing 50 % fresh formamide at a flow rate of 0.5 mL/min. The column was calibrated using commercially available protein standards of known molecular weight (Amersham
15 Pharmacia, Piscataway, New Jersey).

4. Direct ELISA

Greiner high-binding 96 well plates were coated with 100 μ l HPLC fractions overnight at 4°C. Plates were washed three times (5 minutes each) with TBS-Tween and exposed to 100 μ l of 1:2000 primary antibody for 1 hour at room
20 temperature. Plates were washed three times and exposed to 100 μ l of 1:2000 HRP-conjugated goat anti-rabbit IgG. Following a one hour incubation at room temperature, plates were washed again three times with TBS-Tween and color was developed using 3,3', 5,5', tetramethylbenzidine and H₂O₂ for 10 minutes at room temperature. Color development was stopped by the addition of 25 μ L 1N H₂SO₄
25 and analyzed at 450 nm.

5. Immunoprecipitation

Aliquots of normal human serum diluted 1:100 in immunoprecipitation buffer (0.1 M Tris, pH 7.2, 0.15 M NaCl, .05% Tween 20, and 1% aprotinin) were incubated sequentially with 0.1 mL each of (a) Protein G agarose
30 (Kirkegaard & Perry; Gaithersburg, MD), (b) normal rabbit serum IgGs bound to Protein G agarose; (c) rabbit anti-BSP antibodies bound to Protein G agarose.

After incubation for 1 hour at 4°C the reaction was terminated by centrifugation at 10,000 x g for 5 minutes and the supernatant taken to the next immunoprecipitation. The first two incubations removed proteins binding non-specifically to agarose and to normal rabbit serum. The anti-BSP incubation
5 immunoprecipitate was dissolved in gel sample buffer and analyzed by 4 – 20 % gradient SDS-PAGE.

6. Production of Recombinant SIBLINGS Proteins

Adenoviral constructs encoding BSP (rBSP), DMP1 (rDMP1), and OPN (rOPN) were generated by subcloning BSP (Fisher et al., *J. Biol. Chem.*
10 265(4):2347-2351, 1990), BSP-KAE or OPN (Young et al. *Genomics* 7:491-502, 1990; and Kiefer et al., *Nucleic Acid Research* 17:3306, 1989) cDNA into high expression, replication-deficient adenovirus (Ad5) using EF-1 (BSP) and CMV (BSP-KAE, OPN, DMP1) promoters, respectively. The RGD->KAE constructs were made using *in situ* mutagenesis and the entire insert checked for fidelity.
15 Adenoviruses were plaque-selected and propagated on HEK 293 cells (ATCC #CRL1573). Cells were harvested when cytopathic effects were present and lysed by 5 freeze-thaw cycles. Cellular debris was removed and viral particles were purified by banding twice on CsCl. After dialysis in Tris/ MgCl₂ / Glycerol Buffer at 4°C, viruses were aliquoted and frozen at -70°C. Evaluation of viral
20 titers was carried out by plaque formation of virus dilutions on HEK293 cells (Becker et al., *Methods Cell Biol.* 43(Pt A):161-189, 1994). Typically ~2-4 x 10¹¹ pfu/mL were obtained from one viral preparation. Recombinant proteins were generated by infecting subconfluent normal human marrow stromal fibroblasts with 10,000 pfu/cell. Cells were maintained in alpha-MEM, 20 % FBS and 100 IU/mL
25 penicillin/100 µg/mL streptomycin in a humidified atmosphere of 95% air and 5% CO₂ at 37°C. Medium was changed to serum-free conditions after 48 hours. Subsequently, medium was collected every 24-48 hours and frozen at -70°C. Aliquots were assayed by SDS-Gel electrophoresis and Western blot for BSP, DMP1, and OPN expression. Expression was found to be at highest levels at ~168
30 hours post infection. The proteins were purified by routine column chromatography. Native BSP, BSP-KAE, DMP1, and OPN proteins were purified

by diluting medium from normal human marrow stromal fibroblast cells 1:1 with 40 mM phosphate buffer pH 7.4 and loading directly on a 5.0 x 2.0 cm column packed with ToyoPearl TSK QAE resin. A linear salt gradient to 2.0 M NaCl was employed to purify the BSP, DMP1, and OPN to ~95% purity as measured by SDS PAGE.

7. Biotinylated BSP

12 μ g of recombinant human BSP was dissolved in 50 μ L of PBS. 2 μ L of a fresh 1 mg/mL solution of NHS-LC-Biotin (Pierce Chemical Co., Chicago, IL) was added and the reaction incubated at room temperature for 45 min. The unreacted biotin was removed by repeated washing with TBS-Tween and centrifugation in a Microcon 30 (Amicon, Beverly, MA). A final volume of 50 μ L was retained.

8. Alternate Complement Mediated Cell Lysis Assay

Murine erythroleukemia (MEL) cells (a gift of Dr. Marilyn Farquhar, Univ. CA San Diego.) grown in DMEM containing 10 % fetal bovine serum and 4 mM glutamine were rinsed three times with gelatin veronal buffer with Mg_2^+ and EGTA (GVB-MgEGTA, Sigma, St. Louis, MO). Cells were resuspended in GVB-MgEGTA at a density of 5×10^6 cells/mL and incubated at 37°C with different concentrations of normal human serum diluted in GVB-MgEGTA. After 2 h, cells were harvested for trypan blue exclusion assay by removing a 50 μ L aliquot, incubating for 15 minutes in 0.4 % trypan blue, and counting viable cells under an inverted microscope. The thiazolium blue assay was carried out at identical serum dilutions by incubating a 50 μ L aliquot of the cell suspension in an equal volume of 1 mg/mL thiazolyl blue (MTT) for 45 min. Cell viability was determined spectrophotometrically by absorbance at 560 nm. Cells in GVB-MgEGTA buffer were pre-incubated with 10 μ g of either rBSP, rDMP1, or rOPN in 1 mL for 10 minutes at 37°C. Normal human serum collected for good complement activity was then added at a dilution of 1:10 and the cells returned to 37°C for 2 hours and cell viability was determined by trypan blue-exclusion and MTT assays. For the assay of human cancer cell lines, loosely adherent MCF-7 cells were selected for by sequential growth in EMEM with 2 mM L-glutamine

and Earle's balanced salt solution adjusted to contain 1.5 g/L sodium bicarbonate, 0.1 mM non-essential amino acids and 1.0 mM sodium pyruvate and 10% fetal bovine serum, while U-266 cells were cultured in RPMI 1640 medium containing 15 % fetal bovine serum. Cells were collected by centrifugation and rinsed three
5 times with GVB-MgEGTA buffer and subsequently treated exactly as for MEL cells, substituting guinea pig serum (Sigma, St. Louis, MO) for human serum.

EXAMPLES

10

EXAMPLE 1

Isolation of SIBLINGS Proteins and the Relatively Acidic form of BSP

A. Disruption of SIBLINGS/Factor H Complex

A variety of methods of disrupting protein-protein interactions are well known in the art. Such methods can involve treatment with harsh chemicals
15 and/or severe temperatures. Typically protein complexes such as the SIBLINGS/Factor H complex can be disrupted through the use of reducing agents, denaturants and/or increased temperatures. The SIBLINGS/Factor H complex also can be disrupted using a combination of reducing agents, denaturants, and increased temperatures. The particular method of disrupting the complex is not
20 critical, as long as the complex is disrupted.

Reducing agents are agents that are capable of donating hydrogen atoms, and as such serve to disrupt disulfide bonds between amino acids. These agents particularly disrupt disulfide bonds that commonly link two proteins or portions of the same protein together. Commonly used reducing reagents are β -
25 mercaptoethanol, 1,4-dithiothieitol (DTT) and trialkyl phosphines. Additionally, the ability of a reducing agent to facilitate the separation of SIBLINGS proteins from Factor H will be increased by increasing the temperature at which the solution containing the reducing agent and SIBLINGS/Factor H complex is incubated.

30 Denaturants serve to relax the conformational structure of proteins, and there are a variety of denaturants that can be employed to effectuate the separation of SIBLINGS proteins from Factor H. Examples of denaturants that are

particularly useful in conjunction with ion exchange columns are urea and formamide. However, denaturants such as guanidine hydrochloride or guanidine thiocyanate also may be useful for separating Factor H from SIBLINGS proteins in non-ion exchange based methods. The ability of the denaturant to relax the protein
5 will also be enhanced by increasing the temperature at which the sample containing the SIBLINGS/Factor H complex is incubated.

SIBLINGS proteins, such as BSP can also be separated from Factor H by adding competitive binding agents to the sample containing the BSP/Factor H complex. Upon addition to the sample, a competitive binding agent binds to
10 Factor H and displaces BSP. Ideally, the agent used binds to Factor H selectively and not to BSP. Examples of such competitive binding agents are, recombinant fragments of BSP, sialic acids, and synthetic BSP fragments.

Since a variety of methods can be employed to separate Factor H from SIBLINGS proteins, the invention also provides a method of determining when the
15 proteins have become separated. As mentioned above, when Factor H is complexed to SIBLINGS proteins, the relatively acidic BSP, or to the relatively non-acidic BSP, it masks the ionic charge normally associated with protein. Therefore, when the complex is placed on an anion exchange column, it readily flows through the column and the complex fails to bind. Hence, after treatment
20 with one or more denaturants and reducing agents, the sample can be tested for the its ability to bind to an anion exchange column. This assay can confirm whether separation of Factor H from the complex has occurred.

It is also possible to simultaneously confirm that a SIBLINGS protein has been separated from Factor H and measure the SIBLINGS protein level using
25 column chromatography. This can be done by first mixing a sample with an anion exchange resin under conditions that disrupt the Factor H/SIBLINGS complex and then allowing the SIBLINGS protein to bind to the anion exchange resin. The un-complexed SIBLINGS protein will then bind to the resin, denaturants and other contaminants can be washed away, and the SIBLINGS-resin complex can then be
30 directly assayed using SIBLINGS protein specific binding agents.

Additionally, the column conditions can be modified such that either the relatively acidic BSP or the relatively non-acidic BSP remains bound. Thus, BSP specific binding agents used to detect the BSP resin complex will actually be detecting either the relatively non-acidic BSP or the relatively acidic BSP.

5 **B. Separation of the Total Bound SIBLINGS protein from Factor H and the Separation of the Relatively Non-Acidic BSP from the Relatively Acidic BSP**

 The separation of the total bound SIBLINGS protein from Factor H, and the subsequent separation of the relatively acidic and the relatively non-acidic BSP, can be accomplished using multi-step procedures that include various chromatographic techniques (Robyt and White, Biochemical Techniques Theory and Practice, Waveland Press, Inc, 1990). Such multi-step procedures will commonly operate to separate the proteins by exploiting their different physical characteristics. There are many different methods that can be used to effectively separate SIBLINGS protein from Factor H, as well as to separate the relatively acidic BSP from the relatively non-acidic SIBLINGS protein. Therefore, the following discussion provides a broad description of various chromatography techniques that can be used in either a single step process or in multi-step process. Furthermore, it is possible that a single variety of chromatography be repeatedly used to effectuate either the isolation of the relatively acidic SIBLINGS protein, or the entire population of SIBLINGS protein present in the sample.

1. Column Chromatography

 One method of isolating the SIBLINGS protein and/or relatively acidic BSP is through adsorption chromatography. This method involves the exploitation of the differential affinity of the protein for the medium in the column, and the affinity of the protein to the eluting solvent. An example of a suitable medium for use in the column is hydroxyapatite, which is a crystalline form of calcium phosphate. This medium tends to adsorb acidic proteins (such as BSP), which can subsequently be eluted with phosphate ions, which have a high affinity for the calcium ions present in the hydroxyapatite.

 Ion exchange chromatography can also be used to isolate SIBLINGS protein, and/or the relatively acidic BSP. This approach is a variation of

adsorption chromatography, in which the solid adsorbent has charged groups chemically linked to an inert solid. The ionic charge of the SIBLINGS protein then causes the SIBLINGS protein to become linked to the chemically charged groups on the solid support, and the SIBLINGS protein is subsequently released by
5 passing a solution containing an ion gradient over the solid adsorbent. Examples of solid supports that can be used in this technique include DEAE-cellulose, DEAE-Sephadex, DEAE-Bio-Gel, DEAE cellulose, DEAE sepharose, DEAE sephacryl, DEAE trisacryl, Q sepharose, ecteola cellulose, QAE cellulose, express ion exchanger Q, PEI cellulose, and other polystyrene-based anion exchangers.

10 Another type of adsorption chromatography which can be used to isolate SIBLINGS protein and/or the relatively acidic form of BSP is affinity chromatography. This method involves covalently linking to an inert solid support a ligand which has binding affinity for the protein of interest, which in this case is either SIBLINGS protein or the relatively acidic BSP. Commonly, the ligand is a
15 specific binding agent, which selectively binds the protein of interest as it contacts the solid support. Alternatively, the SIBLINGS protein and/or the relatively acidic BSP can be separated based upon hydrophobicity. For example, alkyl chains may be linked to the inert support, to supply sites for hydrophobic bonding interactions. The eluting solvent then contains a hydrophobic gradient.

20 High performance liquid chromatography provides yet another method that can be utilized to isolate SIBLINGS protein and/or the relatively acidic BSP. All of the major classes of chromatographic separations are possible using this method, for example, adsorption, liquid-liquid partition, ion exchange, exclusion, and affinity chromatography can be used in conjunction with HPLC. Additionally, the
25 HPLC method allows for reversed phase and ion pair partition. Reversed phase partition allows for quick elution using a non-polar stationary phase and a polar mobile phase, while ion pair partition involves pairing the charged polar substance with its counter ion to create a less polar species that then flows through the column.

2. Electrophoresis

Electrophoresis is a well-established technique for the separation and analysis of mixtures by differential migration and separation of molecules in an electric field, based on differences in mobility through a support. Many different forms of electrophoresis have been developed to permit the separation of different classes of compounds. These forms include paper and cellulose acetate electrophoresis, thin-layer electrophoresis, gel electrophoresis, immunoelectrophoresis and isoelectric focusing. Paper electrophoresis operates best for the separation of relatively low molecular weight protein molecules, and gel electrophoresis is a much better candidate for use in isolating the relatively acidic BSP. There are a variety of different matrices available for forming gels, but matrices with a relatively smaller pore size are most suitable for the separation of two similar molecules. Furthermore, proteins may be separated based upon size by conducting the electrophoresis under disassociating conditions, for example using SDS. The SDS relaxes the protein and masks the ionic charge of the protein, which leaves protein size as the only distinguishing characteristic. Hence in the case of the relatively acidic BSP, SDS would not be used for separating the two populations of BSP (the relatively acidic versus the relatively non-acidic), because the separation will depend on the ionic charge that would be masked by SDS.

Separation can also be achieved using immunoelectrophoresis, in which proteins are separated on a gel based upon their charge-to-mass ratio and their antigenicity. This is done by first separating the proteins on a gel, and then adding an antibody specific for the protein of interest to a well created in the gel, and allowing the antibody to diffuse through the gel. Precipitate then forms in regions in which the antibody reacts with the protein.

The relatively acidic form of BSP can also be isolated through the use of isoelectric focusing. In this type of electrophoresis the protein is placed on a substrate having a pH gradient, and it will move under the influence of an electrical field until it reaches an equilibrium point with the pH gradient.

Finally, the characterization of the relatively acidic BSP population indicates that other SIBLINGS proteins may exist in multiple forms. Moreover, SIBLINGS proteins may also have relatively acidic population and these populations can be detected using the above described techniques.

5

EXAMPLE 2

Identifying SIBLINGS Proteins with Specific Binding Agents

A SIBLINGS protein can be identified using specific binding agents. These specific binding agents may bind for example, BSP generally, or they may
10 selectively bind to the relatively acidic BSP, the relatively non-acidic BSP, or the exposed portion of SIBLINGS protein in the SIBLINGS/Factor H complex. The latter specific binding agent will be particularly useful for more accurately determining the total amount of SIBLINGS protein in the biological specimen without the need to separate the SIBLINGS protein from the Factor H.

15 A. Antibodies

Antibody preparations prepared according to these protocols are useful in quantitative immunoassays such as ELISAs, Western blotting and RIAs, to determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen
20 in a biological specimen.

1. Production of an Antibody to SIBLINGS Proteins

Monoclonal or polyclonal antibodies may be produced to SIBLINGS proteins or portions thereof. Optimally, antibodies raised against a particular SIBLINGS protein, such as BSP will specifically detect BSP. That is, antibodies
25 raised against BSP would recognize and bind BSP and would not substantially recognize or bind to other proteins found in human cells. The determination that an antibody specifically detects a SIBLINGS protein, such as BSP is made by any one of a number of standard immunoassay methods; for instance, the Western blotting technique (Sambrook et al., In Mol. Clon.: A Lab. Man., Cold Spring
30 Harbor, N.Y., 1989). To determine that a given antibody preparation (such as one produced in a mouse against the human BSP) specifically detects BSP by Western

blotting, serum is extracted from a subject (for example, to obtain lymphocytes) and electrophoresed on a sodium dodecyl sulfate-polyacrylamide gel. The proteins are then transferred to a membrane (for example, nitrocellulose or nylon) by Western blotting, and the antibody preparation is incubated with the membrane.

5 After washing the membrane to remove non-specifically bound antibodies, the presence of specifically bound antibodies is detected by the use of an anti-mouse antibody conjugated to an enzyme such as alkaline phosphatase; application of the substrate 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium results in the production of a dense blue compound by immuno-localized alkaline
10 phosphatase. Antibodies that specifically detect SIBLINGS protein will, by this technique, be shown to bind substantially only the SIBLINGS protein band (which will be localized at a given position on the gel determined by its molecular weight). Non-specific binding of the antibody to other proteins may occur and may be detectable as a weak signal on the Western blot. The non-specific nature
15 of this binding will be recognized by one skilled in the art by the weak signal obtained on the Western blot relative to the strong primary signal arising from the specific antibody-SIBLINGS protein binding.

Antibodies that specifically bind to SIBLINGS protein belong to a class of molecules that are referred to herein as specific binding agents. Specific binding
20 agents that are capable of specifically binding to SIBLINGS protein may include polyclonal antibodies, monoclonal antibodies (including humanized monoclonal antibodies) and fragments of monoclonal antibodies such as Fab, F(ab γ)₂ and Fv fragments, as well as any other agent capable of specifically binding to SIBLINGS proteins.

25 Depending on the desired specificity of the antibody, different forms of a SIBLINGS protein can be used as immunogens for injection into subjects for the creation of antibodies. To create antibodies which recognize the exposed portion of a SIBLINGS protein when it is bound to Factor H, the entire SIBLINGS/Factor H complex can be used as an immunogen. Similarly, if the creation of an antibody
30 specific for the relatively acidic BSP is desired, the relatively acidic BSP can be isolated and used to inoculate a mouse. Additionally, as mentioned below,

fragments of the relatively acidic BSP or fragments of the exposed portion of BSP can be used as immunogens for the creation of specific binding agents.

Regardless of the particular form of SIBLINGS protein used to create the immunogen the concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few micrograms per milliliter. Alternatively, as previously mentioned, peptide fragments of the SIBLINGS proteins may be utilized as immunogens. Such fragments may be chemically synthesized using standard methods, or may be obtained by cleavage of the whole SIBLINGS protein followed by purification of the desired peptide fragments. Peptides as short as 3 or 4 amino acids in length are immunogenic when presented to the immune system in the context of a Major Histocompatibility Complex (MHC) molecule, such as MHC class I or MHC class II. Accordingly, peptides comprising at least 3 and preferably at least 4, 5, 6 or more consecutive amino acids of the disclosed SIBLINGS protein amino acid sequences may be employed as immunogens to raise antibodies.

Because naturally occurring epitopes on proteins frequently include amino acid residues that are not adjacently arranged in the peptide when the peptide sequence is viewed as a linear molecule, it may be advantageous to utilize longer peptide fragments from the SIBLINGS protein amino acid sequence in order to raise antibodies. Thus, for example, peptides that comprise at least 10, 15, 20, 25 or 30 consecutive amino acid residues of the BSP amino acid sequence may be employed. Monoclonal or polyclonal antibodies to the intact BSP or peptide fragments of this protein may be prepared as described below.

2. Monoclonal Antibody Production by Hybridoma Fusion

Monoclonal antibody to epitopes of BSP identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler and Milstein, Nature **256**:495, 1975, or derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein over a period of a few weeks. The mouse is then sacrificed, and the antibody-producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused

cells destroyed by growth of the system on selective media such as aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate, where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the
5 supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, *Enzymol.* 70:419, 1980, and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Harlow and Lane, *Antibodies, A Lab. Man.*, Cold
10 Spring Harbor, N.Y., 1988. In addition, protocols for producing humanized forms of monoclonal antibodies (for therapeutic applications) and fragments of monoclonal antibodies are known in the art.

3. Polyclonal Antibody Production by Immunization

Polyclonal antiserum containing antibodies to heterogeneous epitopes of a
15 single protein can be prepared by immunizing suitable animals with the expressed protein, which can be unmodified or modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than others and may require the use of carriers and adjuvant. Also,
20 host animals vary in response to site of inoculations and dose, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appear to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis et al., *J. Clin. Endocrinol. Metab.* 33:988-991, 1971.

25 Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony et al., In *Handbook of Experim. Immunol.*, Wier, D. ed., ch. 19. Blackwell, 1973. Plateau
30 concentration of antibody is usually in the range of 0.1 to 0.2 mg/mL of serum

(about 12 M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher, 1980.

4. Antibodies Raised by Injection of SIBLINGS protein cDNA

5 Antibodies may also be raised against SIBLINGS proteins by subcutaneous injection of a DNA vector that expresses the SIBLINGS protein in laboratory animals, such as mice. Delivery of the recombinant vector into the animals may be achieved using a hand-held form of the Biolistic system (Sanford et al., *Particulate Sci. Technol.* 5:27-37, 1987) as described by Tang et al., *Nature* 10 (London) 356:152-154, 1992. Expression vectors suitable for this purpose may include those that express a cDNA encoding a SIBLINGS protein under the transcriptional control of either the human β -actin promoter or the cytomegalovirus (CMV) promoter. Methods of administering naked DNA to animals in a manner to cause expression of that DNA in the body of the animal are well known and are 15 described, for example, in U.S. Patent Nos. 5,620,896 (DNA vaccines against rotavirus infection), 5,643,578 (Immunization by inoculation of DNA transcription unit) and 5,593,972 (Genetic immunization), and references cited therein.

5. Antibody Fragments

20 Antibody fragments may be used in place of whole antibodies and may be readily expressed in prokaryotic host cells. Methods of making and using immunologically effective portions of monoclonal antibodies, also referred to as antibody fragments, are well known and include those described in Better and Horowitz, *Advances in Gene Technol.: The Mol. Biol. Of Immune Disease & the Immune Response* (ICSU Short Reports), 1989; Glockshuber et al., *Biochem.* 25 29:1362-1367 (A Comparison of Strategies to Stabilize Immunoglobulin F_v Fragments), 1990; and U.S. Patent Nos. 5,648,237 (Expression of Functional Antibody Fragments), 4,946,778 (Single Polypeptide Chain Binding Molecules), and 5,455,030 (Immunotherapy Using Single Chain Polypeptide Binding Molecules), and references cited therein.

6. Humanized Antibodies

Humanized monoclonal antibodies are preferred in clinical applications. Methods of making humanized monoclonal antibodies are well known, and include those described in U.S. Patent Nos. 5,585,089 (Humanized Immunoglobulins),
5 5,565,332 (Production of Chimeric Antibodies--A Combinatorial Approach),
5,225,539 (Recombinant Altered Antibodies And Methods Of Making Altered Antibodies), 5,693,761 (Polynucleotides Encoding Improved Humanized Immunoglobulins), 5,693,762 (Humanized Immunoglobulins), 5,585,089 (Humanized Immunoglobulins), and 5,530,101 (Humanized Immunoglobulins),
10 and references cited therein.

Antibody preparations prepared according to these protocols are useful in quantitative immunoassays to determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample.

15

EXAMPLE 3

Characterizing SIBLINGS Protein Populations in Biological Specimens

In addition to the assays provided above, the present invention allows for further characterization of SIBLINGS protein populations in biological specimens,
20 and the development of screening assays that can detect tumors, as well as abnormal bone turnover. It is additionally contemplated that tumors such as sarcomas and carcinomas, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon
25 carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumor,
30 cervical cancer, testicular tumor, bladder carcinoma, CNS tumors (such as a glioma, astrocytoma, medulloblastoma, craniopharygioma, ependymoma,

pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma and retinoblastoma) may prove to express BSP and be detectable through the use of the disclosed assays.

A. Development of Screening Assays

5 The invention allows for the development of screening assays that can detect other disease states which are associated with elevated SIBLINGS protein levels, and/or shifts in the population of BSP from the relatively non-acidic BSP to the relatively acidic form of BSP. This can be done by separating the SIBLINGS protein from Factor H and correlating the relative concentration of SIBLINGS
10 protein in the test sample to the relative concentration in a control. The control in many cases is a like sample from a subject that does not have the particular malady in question. For example, a screening assay can be developed to detect abnormal bone turnover in subjects having osteoporosis or osteoarthritis, by taking a sample from the subject that is know to have the particular malady, and characterizing the
15 SIBLINGS protein concentration using the methods provided above. This same procedure can be then be used to characterize a sample from a normal subject and subsequently a correlation can be drawn between the results from the two subjects. If present the correlation can then be used to detect the malady in other subjects.

**B. Controls For Assays In Which The Relatively Acidic BSP Is
20 Used As A Marker**

The above data shows subjects suffering with breast, prostate, thyroid, multiple myeloma and lung tumors have readily distinguishable populations of BSP. These differences have permitted the development of an assay that can be used to detect the presence or recurrence of such tumors (such as recurrence
25 following surgical excision of a tumor). Generally, such assays include a control that allows a comparison to be made between a normal individual and the test subject. However, in some cases this control can be recombinantly expressed BSP, or the test sample can be classified as positive for a tumor based upon the ratio of the relatively non-acidic BSP to the relatively acidic BSP. The above
30 separation methods describe different mechanisms for separating proteins based upon ionic charge and these methods can be used to provide information on the

relative concentration of the relatively non-acidic form of BSP, as well as the relatively acidic form of BSP. Therefore, used in this way the assay does not require the use of an external control; rather the ratio itself can be used to indicate the presence of a tumor.

5

EXAMPLE 4

Use Of Members of the SIBLINGS Family of Proteins to Confer Protection against Complement

The above-described methods and results show that when members of the SIBLINGS family of proteins are associated with cells they can serve to protect the cells from complement mediated lysis. In the case of tumor cells it is likely that the expression of SIBLINGS proteins serves to protect the tumor from complement lysis. This protection may be especially important during the process of metastasis when tumor cells are invading other tissue such as bone.

The complement protective activity of SIBLINGS proteins can also be exploited to protect cells and/or implants from complement mediated immune responses. In the case of implants SIBLINGS proteins can be used to coat implants such as described in U.S. Patent No. 6,024,918 to Hendriks, et al. or they can be delivered from a reservoir that is placed in the body. This protection can be imparted to a cell by either coating the cell with the SIBLINGS protein, or genetically engineering a cell to express the SIBLINGS protein. Used in this way a cell that has been engineered to express a transgene is designed to also express one or more of the members of the SIBLINGS family of proteins. The co-expression of the SIBLINGS protein then facilitates the binding of Factor H to the transgenic cell. This binding results in the cell becoming protected from complement mediated lysis. Expressing the SIBLINGS protein with a membrane localization peptide could further enhance the protection conferred to the transgenic cell.

The complement protective activity of SIBLINGS proteins also could be used to provide protection against non-self cells that are introduced into a subject's body, for example by implantation of a graft. Used in this way a transgene

encoding the SIBLING protein could be delivered to the graft via various vectors such as described below.

A. Expression Cassette Transfer

5 A preferred method of supplying exogenous SIBLINGS proteins is by transferring a vector comprising an expression cassette to cells associated with the region of interest such that the cells express the SIBLINGS protein.

1. Expression Cassettes

Expression cassettes employed in the present inventive methods are of any type appropriate for cells containing the cassette to express the protein of interest.
10 Thus, for example, an expression cassette comprises a polynucleotide encoding BSP operably linked to a promoter.

Any promoter and/or enhancer sequence appropriate for controlling expression of polynucleotides from the vector can be used in constructing an expression cassette. While such promoters/enhancer elements are well known in
15 the art, examples of suitable promoters include prokaryotic promoters or viral promoters, (e.g., ITRs, or LTRs; immediate early viral promoters, such as herpesvirus IE promoters, or cytomegalovirus (CMV) IE promoters; or other viral promoters, such as Rous Sarcoma Virus (RSV) promoters, or Murine Leukemia Virus (MLV) promoters). Other suitable promoters are eukaryotic promoters,
20 such as constitutively active promoters (e.g., the β -actin promoter), signal specific promoters (e.g., inducible promoters, such as a promoter responsive to TNF), or tissue-specific promoters, (e.g., those active in epidermal tissue, dermal tissue, tissue of the digestive organs such as cells of the esophagus, stomach, intestines, or colon), smooth muscles, such as vascular smooth muscles, cardiac muscles,
25 skeletal muscles, lung tissue, hepatocytes, lymphocytes, endothelial cells, sclerocytes, kidney cells, glandular cells (e.g., those in the thymus, ovaries, testicles, pancreas, adrenals, pituitary, etc.), tumor cells, cells in connective tissue, cells in the central nervous system (e.g., neurons, neuralgia, etc.), cells in the peripheral nervous system, or other cells of interest.

30 Which promoter is used in a given expression cassette will depend, in part, on the choice of vector. Thus, for example, an expression cassette can comprise a

native retroviral LTR promoter operably linked to a coding polynucleotide when the vector is a retroviral vector.

In addition to a promoter, an expression cassette will also contain a polynucleotide encoding the protein of interest. Preferably, the polynucleotide is a synthetic DNA, cDNA or genomic DNA fragment encoding the SIBLINGS protein or fragment thereof that imparts protection against complement mediated lysis to the cell that expresses it. Expression cassettes can also include other polynucleotides, such as a polyadenylation sequence. Also, expression cassettes can encode more than one SIBLINGS protein.

10 2. Vectors

An expression cassette for use in the present inventive methods such as those described supra is contained within a vector. Of course, in addition to the SIBLINGS protein encoding sequence, the vector can include other expression cassettes, such as, for example, cassettes for expressing a selectable marker (e.g., β -gal or a marker conferring resistance to a toxin), a pharmacologically active protein, a transcription factor, or other biologically active substances.

Any vector appropriate for transferring an exogenous expression cassette to a cell is included within the scope of the present inventive methods. Preferably, the vector is a viral vector. Examples of viral vectors employed in accordance with the present inventive method include, but are not limited to, retroviral vectors, adenoviral vectors, adeno-associated viral vectors, herpesviral vectors, SV40 viral vectors, polyoma virus vectors, papilloma virus vectors, picornavirus vectors, vaccinia virus vectors, or other suitable vectors.

In addition to viral vectors, any non-viral vector capable of expression upon infection of target cells can be used in the present inventive methods. Preferably, a non-viral vector is a plasmid.

The skilled artisan will be able to incorporate an expression cassette into the nucleic acid sequence of the vector. Methods for incorporating expression cassettes into viral vectors are well known in the art (see e.g., Sambrook, et al. *Molecular Cloning: a Laboratory Manual*, 2d edition, Cold Spring Harbor Press, 1989) and include direct cloning, site specific recombination using recombinases,

such as the flp recombinase or the cre-lox recombinase system (reviewed in Kilby et al., Trends in Genetics, 9:413-21, 1993), homologous recombination, and other suitable methods of constructing a recombinant vector.

3. Target Cells

5 The present inventive methods also comprise transferring an expression cassette to cells associated with the region of interest (i.e., a graft). Any suitable cells associated with the region of interest which are capable of supporting expression of the coding polynucleotide of the cassette are encompassed within the present inventive methods. Such cells can be cells *in situ*, such as cells of the graft
10 tissue.

Following transfer of the vector comprising the expression cassette to the cells *in vitro*, the cells associated with the region are transferred to the region of interest. Transfer of cells *in vitro* to a region of interest is accomplished by known means. For example, cells can be transferred to internal tissue of a patient by
15 methods described herein. Further methods of transferring cells containing an expression cassette are described in International Patent Application No. WO 96/00006, (Billiar et al.).

4. Vector Transfer

20 The present inventive methods comprise transferring a vector comprising an expression cassette, such as those described herein, to cells associated with the region of interest (e.g., graft). Any method of transferring the expression cassette to the cells is appropriate so long as a product of the expression cassette is produced in the cells.

a. Non-Viral Vectors

25 Any means of introducing non-viral vectors into target cells, such as by direct DNA injection, electroporation, calcium-phosphate mediated cell transfection, lipofectamine, DAEA-dextran-mediated cell transfection, polybrene-mediated delivery, host cell fusion, microinjection, and polylysine-mediated cell transfection is appropriate within the present inventive context. Such methods are
30 well known in the art, and some are described in International Patent Application No. WO 96/00006, (Billiar et al.). A preferred method of transferring the

expression cassette within a nonviral vector, such as a plasmid, is via liposome-mediated transfection of the target cells, such as endothelial cells, *in vitro* or *in situ*. If the vector is transfected *in vitro*, the cells associated with the region of interest are subsequently transferred to the patient.

5

b. Viral Vectors

Transfer of an expression cassette to cells associated with the region of interest where the cassette is within a viral vector is accomplished by infecting the cells with the virus. Where the virus is a retrovirus, the virus can be first transfected into an appropriate packaging cell line for generation of infectious virus. Cells associated with the region of interest can be infected *in vitro* or *in situ*. Moreover, the mode of transfer of the expression cassette does not depend on the identity of the coding polynucleotide.

10

c. In vitro Delivery

Infectious viruses are used to infect cultured cells *in vitro*. The precise method for infecting target cells *in vitro* will depend upon the viral vector used; however, such methods are well known in the art. Some suitable methods are described in International Patent Application No. WO 96/00006, (Billiar et al.).

15

5. In situ Delivery

Some applications of the present inventive methods involve transfer of expression cassettes *in situ*. *In situ* transfer can be either *in vivo* (i.e., to a wound or to a graft following implantation), or *ex vivo* (i.e., to a graft prior to implantation). Any method of delivering the vector comprising the expression cassette to cells associated with the region of interest *in situ* is within the present methods. Such methods also apply to transfer of exogenous cells to the region of interest. Methods for *in situ* delivery of vectors preferably involve physically segregating the region of interest from the remainder of the patient's tissue in order to properly target the vector within the region. Upon segregation, the vector is applied to the region of interest in a manner appropriate to transfer the expression cassette into the cells associated with the region of interest.

20

25

30

Tissue *ex vivo*, such as, for example, a graft, is completely isolated from the patient. For *ex vivo* vector transfer, the vector can be applied to the tissue in

any suitable manner, e.g., in a carrier appropriate for transferring the vector. For example, the tissue can be incubated in a carrier containing the vector particles by any suitable incubation method. Other tissue can be perfused with the solution containing the vector.

5 The vectors or exogenous cells preferably remain in contact with the wound or tissue for a period of time sufficient to promote the transfer. For example, a carrier comprising a vector will remain in contact with the wound or *ex vivo* tissue from about one minute to about 2 hours, more preferably between 10 minutes and an hour, and most preferably from about 20 minutes to 40 minutes. In many
10 applications, the carrier will optimally remain in contact with the wound or tissue for about 30 minutes.

6. Internal Delivery

For *in situ* delivery of a vector internally, the region of interest desirably is further segregated from the remainder of the subject's tissue. Any of a variety of
15 known surgical procedures for physically segregating the region of interest is appropriate. Various endovascular surgical techniques appropriate for segregating a region of interest are available, depending upon the location of the target.

Endovascular surgical procedures include, but are not limited to, balloon angioplasty, intravascular stents, laser-assisted balloon angioplasty, double balloon
20 catheterization, mechanical endarterectomy and vascular endoscopy. For a review of endovascular alternatives, see generally Ahn, "Endovascular Surgery," in Vascular Surgery, A Comprehensive Review, Ed. W. S. Moore, W. B. Saunders & Co., Philadelphia, 1993.

Several catheter designs can be utilized for local delivery of a vector. One
25 catheter design consists of two independently inflated balloons, one proximal and one distal to the vascular delivery site. Inflation of these balloons provides an evacuated isolated arterial segment into which vectors for expression cassette delivery can be infused. This system is, however, limited by absence of distal arterial perfusion. A second catheter design developed by Wolinsky allows the
30 infusion of the SIBLINGS protein-encoding vector through 25-100 μ M pores under pressures up to 5 atm. This perfusion pressure increases the depth of penetration

by the vectors and additionally increases expression cassette transfer efficiency. Yet another catheter design utilizes an expandable stent which traps the balloon against the arterial wall and allows intramural delivery of the expression cassette through spaces in the stent material. Additionally, these stents can be modified
5 with burrs that create holes deeper in the vessel wall and allow flow of the expression cassette delivery agents to these sites to allow more uniform delivery of the expression cassette throughout the vessel wall.

Another delivery mechanism is to coat the catheter with a hydrophilic polyacrylic acid polymer that acts as a drug-absorbing sponge. By disrupting the
10 vessel during the angioplasty procedure, this hydrogel is deposited within the vessel wall and will allow sustained delivery of the vector at the arterial wound site. Additionally, the iontophoretic balloon catheter is a catheter design that uses low electrical current to change the cell membrane polarity and allow the diffusion of charged DNA particles into the cell. This is a potential delivery mechanism for
15 plasmid DNA expression cassette constructs. Also, biodegradable stents formed from agents such as an ethylenevinyl acetic copolymer are appropriate for localized delivery to vascular tissue. Alternatively, an intravascular stent can be utilized wherein the endovascular scaffold of the stent is bathed in a ointment, cream, lotion, colloidal dispersion such as a gel or magma or any other acceptable carrier
20 which comprises the SIBLINGS protein encoding vector for delivery to the targeted portion of a vessel segment. This solution is applicable to either an *in situ* or *ex vivo* based vessel delivery.

EXAMPLE 5

25 Additional Examples of Methods for Detecting and/or Quantifying SIBLINGS proteins

The following assays describe the detection of BSP, however, these assays, with appropriate modifications (i.e., antibody substitutions), can also be used to detect other SIBLINGS proteins.

A. Chemiluminescent Sandwich Assay for BSP in Human Serum

Pierce Reacti-Bind White Opaque 96-Well EIA plates (product # 15042) are coated with 100 μ L of 1 μ g/mL Goat anti Mouse IgG (H & L), human serum adsorbed (Kirkegaard & Perry, Gaithersburg, MD) in PBS. The plates are then covered and incubated overnight at 4°C. The next morning the wells are emptied and 250 μ L of 1 % bovine serum albumin is added for 15 min. The albumin is then removed and the plates are washed 3 times with ~300 μ L PBS-0.05 % Tween 20 (PBS-T) for 5 minutes each at room temperature. 100 μ L 1:1000 LFMb-2 (monoclonal to human BSP) in PBS-T is added to each well and the plate is incubated again for 1 hour at room temp and then washed three times (PBS-Tween).

The samples are prepared from 25 μ L human. 50 μ L fresh formamide and 1 μ L of 0.1 M Dithiothreitol is added to the sample and the sample is then heated at 100°C for 5 minutes and diluted to 1 mL with PBS-T containing 0.03 % H_2O_2 . This addition dilutes the fomamide to levels sufficiently low enough to promote normal antibody binding and the H_2O_2 inactivates the remaining DTT.

The BSP standard (5 ng to 0.15 ng) and a negative control (without BSP) are prepared in PBS-T and added to individual wells. 100 μ L diluted sample is also added to the individual wells. The samples and the standards are then incubated 1 hour at room temperature and then washed 3 times with PBS-Tween. 100 μ L of 1:1000 LF-100 polyclonal (rabbit anti-human BSP) antibody in PBS-T plus 1 % normal mouse serum is then added to each well and the plate is incubated for 1 hour at room temperature and then washed 3 times. 100 μ L of 1:50,000 diluted peroxidase-conjugated goat anti rabbit IgG (H&L), multi-serum adsorbed (Kirkegaard & Perry) is then added to each well and the plate is incubated for 1 hour at room temperature, washed 3 times (PBS-Tween).

The results are then visualized by adding 100 μ L of Pierce Supersignal ELISA Pico Chemiluminescent Substrate (product # 37070) and incubating 10 minutes at room temperature. Luminescence can then be detected in a luminometer.

B. Western Blot and Quantitative Immunoanalysis

5 μ L of serum is added to 95 μ L of sample buffer (0.0625 M Tris, pH 6.8, 2.3% SDS, 5 % 2-mercaptoethanol or 1 mM dithiothreitol, 8 M urea or 50% formamide, bromophenol blue tracking dye), the sample is then heated for 5 minutes at 100°C. 10 μ L of the sample is then added to each well of a 4-20 % or 4-12 % gradient acrylamide gel (Novex Corp) and the gel is electrophoresed with 100-200 V constant voltage until the dye front reaches the bottom of the gel. Molecular weight standards (prestained) and BSP standards are also
10 electrophoresed.

The protein is then transferred to nitrocellulose or other western blot membranes under standard conditions. If nitrocellulose is used transfer can be achieved in 60 minutes at 100 volts with cooling in the Novex XCell II Blot Module in Tris/Glycine buffer containing 20% methanol.

15 Protein is then detected by blocking the membrane with 3% BSA-NGS in PBS for 15 minutes and then adding 1% normal goat serum for 45 minutes. An antibody, for instance the polyclonal antibody LF-100, or monoclonal LF-Mb-11 is then diluted at 1:2000 and added to the membrane for 1 hour at room temperature. The membrane is then washed 3 times with PBS-Tween 20 (0.05%).
20 3% BSA-1% NGS is added and then HRP-conjugated second antibody (goat anti-rabbit or goat anti-mouse IgG respectively) at is added to a final concentration of 1:50,000. The membrane is then incubated for 1 hour at room temperature and washed 3 times. Finally, Pierce SuperSignal West Pico Chemiluminescent reagent is added for 5 minutes and the membrane is placed in clear plastic bag and exposed
25 to a chilled digital camera (Fuji LAS-100).

In view of the many possible embodiments to which the principles of the invention may be applied, it should be recognized that the illustrated embodiments are only particular examples of the invention and should not be taken as a limitation on the scope of the invention. Rather, the scope of
30 the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

We Claim:

1. A method of detecting at least one member of the SIBLINGS family of proteins comprising detecting the SIBLINGS protein in a manner that Factor H
5 does not inhibit the detection of the SIBLINGS protein.
2. The method of claim 1 wherein the SIBLINGS protein is in a sample.
3. The method of claim 2 wherein the method further comprises:
10 separating Factor H from the SIBLINGS protein; and
detecting the SIBLINGS protein.
4. The method of claim 3 wherein separating Factor H from the SIBLINGS protein comprises heating the biological sample in the presence of a
15 reducing agent.
5. The method of claim 4 wherein the reducing agent is selected from the group consisting of DTT (1,4-dithiothreitol) and β -mercaptoethanol.
- 20 6. The method of claim 3 wherein the SIBLINGS protein is BSP.
7. The method of claim 6 further comprising separating a relatively acidic BSP from a relatively non-acidic BSP.
- 25 8. The method of claim 7 wherein the relatively acidic BSP is separated from the relatively non-acidic BSP by reacting the BSP with a specific binding agent.
- 30 9. The method of claim 3 wherein the Factor H is separated from the SIBLINGS protein based upon an ionic charge of the SIBLINGS protein and Factor H.

10. The method of claim 3 wherein the Factor H is separated from the SIBLINGS protein by reacting a SIBLINGS/Factor H complex with a specific binding agent specific for the SIBLINGS protein.

5

11. The method of claim 10 wherein the specific binding agent comprises an antibody specific for a portion of the SIBLINGS protein that is exposed in the SIBLINGS/Factor H complex.

10

12. The method of claim 11 wherein the specific binding agent separates and identifies the SIBLINGS protein simultaneously.

15

13. The method of claim 2 wherein the sample is from a subject that is being screened for a presence or a recurrence of a tumor.

15

14. The method of claim 13 wherein the tumor produces a relatively acidic BSP, the method comprising detecting an abnormal presence of the relatively acidic BSP in the sample from the subject.

20

15. A composition comprising a specific binding agent which specifically binds at least one SIBLING protein when the SIBLINGS protein is complexed with Factor H.

25

16. The composition of claim 15 wherein the specific binding agent is a monoclonal antibody.

30

17. The composition of claim 16 wherein the monoclonal antibody specifically recognizes the exposed portion of the SIBLINGS protein when the SIBLINGS protein is complexed with Factor H.

30

18. A hybridoma which expresses the monoclonal antibody of claim 17.

19. A method of detecting a presence of a tumor in a subject from whom a sample is obtained, wherein the tumor produces a relatively acidic BSP, the method comprising:

5 separating Factor H from a Factor H/BSP complex in the sample; and
detecting an abnormal concentration of the relatively acidic BSP

20. A method of making an antibody for detecting a SIBLINGS/Factor H complex, comprising presenting an antigen which is unique to a
10 SIBLINGS/Factor H complex to a subject.

21. An antibody made by the method of claim 20.

22. A transformed organism that expresses a relatively acidic BSP,
15 comprising a cell transformed with a BSP encoding construct that expresses a relatively acidic BSP.

23. The transformed organism of claim 22 wherein the BSP encoding construct comprises a DNA sequence shown in SEQ ID NO: 8.
20

24. A purified relatively acidic BSP produced by an organism of claim 23.

25. A method of conferring protection from a complement mediated
25 immune response to a subject, the method comprising providing within a body a supply of a protective amount of a protein selected from the group consisting of a SIBLINGS protein and a biologically active fragment of a SIBLINGS protein, sufficient to protect the subject from a complement immune response by association with the protein.

30

26. A method of conferring protection from complement mediated lysis to a cell, the method comprising:

providing a vector encoding at least one biologically active SIBLINGS protein;

5 transforming the cell with the vector, wherein the resulting transformed cell expresses the biologically active SIBLINGS protein, and such expression confers resistance to complement mediated lysis to the cell.

27. The method of claim 2, 25, or 26, wherein the SIBLINGS protein is
10 BSP.

28. The method of claim 2, 25, or 26, wherein the SIBLINGS protein is OPN.

29. The method of claim 2, 25, or 26, wherein the SIBLINGS protein is
15 DMP1.

30. The method of claim 2, 25, or 26, wherein the SIBLINGS protein is
20 DSPP.

31. The method of claim 26 wherein the cell is a cell found in a graft.

32. The method of claim 26 wherein the cell is bone marrow cell.

25 33. The method of claim 2 wherein the sample is from a subject suspected of having abnormal bone turnover.

34. The method of claim 33 wherein the subject is suspected of having
osteoporosis.

30

36. The method of claims 2 or 19, wherein the sample comprises a body fluid.

37. The method of claim 35 wherein the body fluid comprises blood of
5 the subject.

38. The method of claim 2 or 19, wherein the tumor is selected from the group consisting of breast, lung, thyroid, multiple myeloma, and prostate.

10 39. The method of claim 14 or 19, further comprising comparing a relative concentration of the relatively acidic BSP in the body fluid to a relative concentration of the relatively acidic BSP in a control sample.

40. The method of claim 14 or 19, wherein detecting the relatively
15 acidic BSP comprises detection by a specific binding agent.

41. The method of claim 40 wherein the specific binding agent comprises a polyclonal antibody.

20 42. The method of claim 40 wherein the specific binding agent comprises a BSP specific monoclonal antibody selected from the group consisting of a monoclonal antibody specific for the relatively acidic BSP, a monoclonal antibody specific for the relatively non-acidic BSP or a monoclonal antibody specific for the exposed portion of BSP when BSP is complexed with Factor H.

25

43. The method of claim 14 or 19, wherein detecting the relatively acidic BSP comprises detection based upon the acidity of an relatively acidic BSP.

44. The method of claim 14 or 19, further comprising detecting the
30 relatively non-acidic form of BSP in the sample and comparing the amount of

relatively non-acidic BSP in the sample to the amount of relatively acidic BSP in the sample.

45. The method of claim 25 wherein providing the supply comprises
5 providing an implant which dispenses the protein, or a cell which expresses the protein.

1/11




FIG. 1A

FIG. 1B

FIG. 1C

FIG. 1

2/11





FIG. 2A

FIG. 2B

FIG. 2C

FIG. 2

3/11



FIGs. 3A and 3B




FIG. 4








FIG. 5

6/11




FIG. 6

7/11




FIG. 7

8/11





FIG. 8

9/11

**FIG. 9**

10/11




Figure 8.

FIG. 10

11/11




FIG. 11A

FIG. 11B

FIG. 11

SEQUENCE LISTING

<110> Fisher et al., Larry

<120> Complex Formed by Small Integrin-Binding Ligand,
N-Linked Glycoproteins (SIBLINGS) and Factor H

<130> 54101

<140>

<141>

<150> 60/128,468

<151> 1999-04-09

<160> 18

<170> PatentIn Ver. 2.1

<210> 1

<211> 317

<212> PRT

<213> Homo sapiens

<400> 1

Met Lys Thr Ala Leu Ile Leu Leu Ser Ile Leu Gly Met Ala Cys Ala
1 5 10 15

Phe Ser Met Lys Asn Leu His Arg Arg Val Lys Ile Glu Asp Ser Glu
20 25 30

Glu Asn Gly Val Phe Lys Tyr Arg Pro Arg Tyr Tyr Leu Tyr Lys His
35 40 45

Ala Tyr Phe Tyr Pro His Leu Lys Arg Phe Pro Val Gln Gly Ser Ser
50 55 60

Asp Ser Ser Glu Glu Asn Gly Asp Asp Ser Ser Glu Glu Glu Glu Glu
65 70 75 80

Glu Glu Glu Thr Ser Asn Glu Gly Glu Asn Asn Glu Glu Ser Asn Glu
85 90 95

Asp Glu Asp Ser Glu Ala Glu Asn Thr Thr Leu Ser Ala Thr Thr Leu
100 105 110

Gly Tyr Gly Glu Asp Ala Thr Pro Gly Thr Gly Tyr Thr Gly Leu Ala
115 120 125

Ala Ile Gln Leu Pro Lys Lys Ala Gly Asp Ile Thr Asn Lys Ala Thr
130 135 140

Lys Glu Lys Glu Ser Asp Glu Glu Glu Glu Glu Glu Glu Glu Gly Asn
145 150 155 160

Glu Asn Glu Glu Ser Glu Ala Glu Val Asp Glu Asn Glu Gln Gly Ile
165 170 175

Asn Gly Thr Ser Thr Asn Ser Thr Glu Ala Glu Asn Gly Asn Gly Ser
180 185 190

Ser Gly Gly Asp Asn Gly Glu Glu Gly Glu Glu Glu Ser Val Thr Gly
 195 200 205
 Ala Asn Ala Glu Gly Thr Thr Glu Thr Gly Gly Gln Gly Lys Gly Thr
 210 215 220
 Ser Lys Thr Thr Thr Ser Pro Asn Gly Gly Phe Glu Pro Thr Thr Pro
 225 230 235 240
 Pro Gln Val Tyr Arg Thr Thr Ser Pro Pro Phe Gly Lys Thr Thr Thr
 245 250 255
 Val Glu Tyr Glu Gly Glu Tyr Glu Tyr Thr Gly Val Asn Glu Tyr Asp
 260 265 270
 Asn Gly Tyr Glu Ile Tyr Glu Ser Glu Asn Gly Glu Pro Arg Gly Asp
 275 280 285
 Asn Tyr Arg Ala Tyr Glu Asp Glu Tyr Ser Tyr Phe Lys Gly Gln Gly
 290 295 300
 Tyr Asp Gly Tyr Asp Gly Gln Asn Tyr Tyr His His Gln
 305 310 315

<210> 2
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:immunogen

<400> 2
 Tyr Glu Ser Glu Asn Gly Glu Pro Arg Gly Asp Asn Tyr Arg Ala Tyr
 1 5 10 15

Glu Asp

<210> 3
 <211> 12
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:immunogen

<400> 3
 Ala Ile Gln Leu Pro Lys Lys Ala Gly Asp Ile Cys
 1 5 10

<210> 4
 <211> 60
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:immunogen

<400> 4

Glu Tyr Glu Gly Glu Tyr Glu Tyr Thr Gly Val Asn Glu Tyr Asp Asn
 1 5 10 15

Gly Tyr Glu Ile Tyr Glu Ser Glu Asn Gly Glu Pro Arg Gly Asp Asn
 20 25 30

Tyr Arg Ala Tyr Glu Asp Glu Tyr Ser Tyr Phe Lys Gly Gln Gly Tyr
 35 40 45

Asp Gly Tyr Asp Gly Gln Asn Tyr Tyr His His Gln
 50 55 60

<210> 5

<211> 152

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:immunogen

<400> 5

Ile Gln Leu Pro Lys Lys Ala Gly Asp Ile Thr Asn Lys Ala Thr Lys
 1 5 10 15

Glu Lys Glu Ser Asp Glu Glu Glu Glu Glu Glu Glu Gly Asn Glu
 20 25 30

Asn Glu Glu Ser Glu Ala Glu Val Asp Glu Asn Glu Gln Gly Ile Asn
 35 40 45

Gly Thr Ser Thr Asn Ser Thr Glu Ala Glu Asn Gly Asn Gly Ser Ser
 50 55 60

Gly Gly Asp Asn Gly Glu Glu Gly Glu Glu Glu Ser Val Thr Gly Ala
 65 70 75 80

Asn Ala Glu Gly Thr Thr Glu Thr Gly Gly Gln Gly Lys Gly Thr Ser
 85 90 95

Lys Thr Thr Thr Ser Pro Asn Gly Gly Phe Glu Pro Thr Thr Pro Pro
 100 105 110

Gln Val Tyr Arg Thr Thr Ser Pro Pro Phe Gly Lys Thr Thr Thr Val
 115 120 125

Glu Tyr Glu Gly Glu Tyr Glu Tyr Thr Gly Val Asn Glu Tyr Asp Asn
 130 135 140

Gly Tyr Glu Ile Tyr Glu Ser Glu
 145 150

<210> 6

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:immunogen

<400> 6

Val Phe Lys Tyr Arg Pro Arg Tyr Tyr Leu Tyr Lys His Ala Tyr Phe
1 5 10 15

Tyr Pro His Leu Lys Arg Phe Pro Val
20 25

<210> 7

<211> 28

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:immunogen

<220>

<221> VARIANT

<222> (5)

<223> Sulfated

<220>

<221> VARIANT

<222> (9)

<223> sulfated

<220>

<221> VARIANT

<222> (11)

<223> sulfated

<220>

<221> VARIANT

<222> (17)

<223> Sulfated

<220>

<221> VARIANT

<222> (21)

<223> sulfated

<220>

<221> VARIANT

<222> (24)

<223> sulfated

<400> 7

Cys Thr Val Glu Tyr Glu Gly Glu Tyr Glu Tyr Thr Gly Ala Asn Asp
1 5 10 15

Tyr Asp Asn Gly Tyr Glu Ile Tyr Glu Ser Glu Asn
20 25

<210> 8

<211> 1037

<212> DNA

<213> Homo sapiens

<400> 8

```

ccttctctgc cctctcactc ccttgagcct gcttcctcac tccaggactg ccagaggaag 60
caatcaccaa aatgaagact gctttaattt tgctcagcat tttgggaatg gcctgtgctt 120
tctcaatgaa aaatttgcat cgaagagtca aaatagagga ttctgaagaa aatggggtct 180
ttaagtacag gccacgatat tatctttaca agcatgccta cttttatcct catttaaaac 240
gatttccagt tcagggcagt agtgactcat ccgaagaaaa tggagatgac agttcagaag 300
aggaggagga agaagaggag acttcaaatg aaggagaaaa caatgaagaa tcgaatgaag 360
atgaagactc tgaggctgag aataccacac tttctgctac aacactgggc tatggagagg 420
acgccacgcc tggcacaggg tatacagggg tagctgcaat ccagcttccc aagaaggctg 480
gggatataac aaacaaagct acaaaagaga aggaaagtga tgaagaagaa gaggaggaag 540
aggaaggaaa tgaaaacgaa gaaagcgaag cagaagtgga tgaaaacgaa caaggcataa 600
acggcaccag taccaacagc acagaggcag aaaacggcaa cggcagcagc ggaggagaca 660
atggagaaga aggggaagaa gaaagtgtca ctggagccaa tgcagaaggc accacagaga 720
ccggagggca gggcaagggc acctcgaaga caacaacctc tccaaatggg ggggttgaac 780
ctacaacccc accacaagtc tatagaacca cttccccacc ttttgggaaa accaccaccg 840
ttgaatacga gggggagtac gaatacacgg gcgtcaatga atacgacaat ggatatgaaa 900
tctatgaaag tgagaacggg gaacctcgtg gggacaatta ccgagcctat gaagatgagt 960
acagctactt taaaggacaa ggctacgatg gctatgatgg tcagaattac taccaccacc 1020
agtgaagctc cagcctg                                     1037

```

<210> 9

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<221> THIOETH

<222> (2)..(17)

<220>

<223> Description of Artificial Sequence: Immunogen

<400> 9

```

Glu Tyr Glu Tyr Thr Gly Val Asn Glu Tyr Asp Asn Gly Tyr Glu Ile
 1             5             10             15

```

```

Tyr Glu Ser Glu Asn Gly Glu Pro
          20

```

<210> 10

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Competitor peptide

<400> 10

```

Gly Arg Gly Asp Ser
 1             5

```

<210> 11

<211> 1447

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (109)..(1011)

<400> 11

gaggcagcag cagcaggagg aggcagacac agcatcgctcg ggaccagact cgtctcaggc 60

cagttgcagc cttctcagcc aaacgccgac caaggaaaac tcactacc atg aga att 117
 Met Arg Ile
 1

gca gtg att tgc ttt tgc ctc cta ggc atc acc tgt gcc ata cca gtt 165
 Ala Val Ile Cys Phe Cys Leu Leu Gly Ile Thr Cys Ala Ile Pro Val
 5 10 15

aaa cag gct gat tct gga agt tct gag gaa aag cag ctt tac aac aaa 213
 Lys Gln Ala Asp Ser Gly Ser Ser Glu Glu Lys Gln Leu Tyr Asn Lys
 20 25 30 35

tac cca gat gct gtg gcc aca tgg cta aac cct gac cca tct cag aag 261
 Tyr Pro Asp Ala Val Ala Thr Trp Leu Asn Pro Asp Pro Ser Gln Lys
 40 45 50

cag aat ctc cta gcc cca cag acc ctt cca agt aag tcc aac gaa agc 309
 Gln Asn Leu Leu Ala Pro Gln Thr Leu Pro Ser Lys Ser Asn Glu Ser
 55 60 65

cat gac cac atg gat gat atg gat gat gaa gat gat gat gac cat gtg 357
 His Asp His Met Asp Asp Met Asp Asp Glu Asp Asp Asp Asp His Val
 70 75 80

gac agc cag gac tcc att gac tcg aac gac tct gat gat gta gat gac 405
 Asp Ser Gln Asp Ser Ile Asp Ser Asn Asp Ser Asp Asp Val Asp Asp
 85 90 95

act gat gat tct cac cag tct gat gag tct cac cat tct gat gaa tct 453
 Thr Asp Asp Ser His Gln Ser Asp Glu Ser His His Ser Asp Glu Ser
 100 105 110 115

gat gaa ctg gtc act gat ttt ccc acg gac ctg cca gca acc gaa gtt 501
 Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala Thr Glu Val
 120 125 130

ttc act cca gtt gtc ccc aca gta gac aca tat gat ggc cga ggt gat 549
 Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly Arg Gly Asp
 135 140 145

agt gtg gtt tat gga ctg agg tca aaa tct aag aag ttt cgc aga cct 597
 Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe Arg Arg Pro
 150 155 160

gac atc cag tac cct gat gct aca gac gag gac atc acc tca cac atg 645
 Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr Ser His Met
 165 170 175

gaa agc gag gag ttg aat ggt gca tac aag gcc atc ccc gtt gcc cag 693
 Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro Val Ala Gln
 180 185 190 195

gac ctg aac gcg cct tct gat tgg gac agc cgt ggg aag gac agt tat 741
 Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys Asp Ser Tyr
 200 205 210

```

gaa acg agt cag ctg gat gac cag agt gct gaa acc cac agc cac aag 789
Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His Ser His Lys
      215                      220                      225

cag tcc aga tta tat aag cgg aaa gcc aat gat gag agc aat gag cat 837
Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser Asn Glu His
      230                      235                      240

tcc gat gtg att gat agt cag gaa ctt tcc aaa gtc agc cgt gaa ttc 885
Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser Arg Glu Phe
      245                      250                      255

cac agc cat gaa ttt cac agc cat gaa gat atg ctg gtt gta gac ccc 933
His Ser His Glu Phe His Ser His Glu Asp Met Leu Val Val Asp Pro
      260                      265                      270                      275

aaa agt aag gaa gaa gat aaa cac ctg aaa ttt cgt att tct cat gaa 981
Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile Ser His Glu
      280                      285                      290

tta gat agt gca tct tct gag gtc aat taa aaggagaaaa aatacaattt 1031
Leu Asp Ser Ala Ser Ser Glu Val Asn
      295                      300

ctcactttgc atttagtcaa aagaaaaaat gctttatagc aaaatgaaag agaacatgaa 1091

atgctttcttt ctcagtttat tgggttgaatg tgtatctatt tgagtctgga aataactaat 1151

gtgtttgata attagtttag tttgtggcct catggaaact ccctgtaaac taaaagcttc 1211

agggttatgt ctatgttcat tctatagaag aaatgcaaac tatcactgta ttttaatat 1271

tgttattctc tcatgaatag aaatttatgt agaagcaaac aaaatacttt taccactta 1331

aaaagagaat ataacatttt atgtcactat aatcttttgt tttttaagtt agtgtatat 1391

ttgttgatgat tatctttttg tgggtgtgaat aaatctttta tcttgaatgt aataag 1447

<210> 12
<211> 300
<212> PRT
<213> Homo sapiens

<400> 12
Met Arg Ile Ala Val Ile Cys Phe Cys Leu Leu Gly Ile Thr Cys Ala
  1          5          10          15
Ile Pro Val Lys Gln Ala Asp Ser Gly Ser Ser Glu Glu Lys Gln Leu
      20          25          30
Tyr Asn Lys Tyr Pro Asp Ala Val Ala Thr Trp Leu Asn Pro Asp Pro
      35          40          45
Ser Gln Lys Gln Asn Leu Leu Ala Pro Gln Thr Leu Pro Ser Lys Ser
      50          55          60
Asn Glu Ser His Asp His Met Asp Asp Met Asp Asp Glu Asp Asp Asp
      65          70          75          80
Asp His Val Asp Ser Gln Asp Ser Ile Asp Ser Asn Asp Ser Asp Asp
      85          90          95
Val Asp Asp Thr Asp Asp Ser His Gln Ser Asp Glu Ser His His Ser
      100          105          110
Asp Glu Ser Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala

```

115	120	125
Thr Glu Val Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly		
130	135	140
Arg Gly Asp Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe		
145	150	155
Arg Arg Pro Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr		160
	165	170
Ser His Met Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro		175
	180	185
Val Ala Gln Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys		190
	195	200
Asp Ser Tyr Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His		205
	210	215
Ser His Lys Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser		220
225	230	235
Asn Glu His Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser		240
	245	250
Arg Glu Phe His Ser His Glu Phe His Ser His Glu Asp Met Leu Val		255
	260	265
Val Asp Pro Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile		270
	275	280
Ser His Glu Leu Asp Ser Ala Ser Ser Glu Val Asn		285
290	295	300

<210> 13
 <211> 1424
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (68)..(1012)

<400> 13
 gaccagactc gtctcaggcc agttgcagcc ttctcagcca aacgccgacc aaggaaaact 60
 cactacc atg aga att gca gtg att tgc ttt tgc ctc cta ggc atc acc 109
 Met Arg Ile Ala Val Ile Cys Phe Cys Leu Leu Gly Ile Thr
 1 5 10
 tgt gcc ata cca gtt aaa cag gct gat tct gga agt tct gag gaa aag 157
 Cys Ala Ile Pro Val Lys Gln Ala Asp Ser Gly Ser Ser Glu Glu Lys
 15 20 25 30
 cag ctt tac aac aaa tac cca gat gct gtg gcc aca tgg cta aac cct 205
 Gln Leu Tyr Asn Lys Tyr Pro Asp Ala Val Ala Thr Trp Leu Asn Pro
 35 40 45
 gac cca tct cag aag cag aat ctc cta gcc cca cag aat gct gtg tcc 253
 Asp Pro Ser Gln Lys Gln Asn Leu Leu Ala Pro Gln Asn Ala Val Ser
 50 55 60
 tct gaa gaa acc aat gac ttt aaa caa gag acc ctt cca agt aag tcc 301
 Ser Glu Glu Thr Asn Asp Phe Lys Gln Glu Thr Leu Pro Ser Lys Ser
 65 70 75
 aac gaa agc cat gac cac atg gat gat atg gat gat gaa gat gat gat 349
 Asn Glu Ser His Asp His Met Asp Asp Met Asp Asp Glu Asp Asp Asp
 80 85 90

gac cat gtg gac agc cag gac tcc att gac tcg aac gac tct gat gat	397
Asp His Val Asp Ser Gln Asp Ser Ile Asp Ser Asn Asp Ser Asp Asp	
95 100 105 110	
gta gat gac act gat gat tct cac cag tct gat gag tct cac cat tct	445
Val Asp Asp Thr Asp Asp Ser His Gln Ser Asp Glu Ser His His Ser	
115 120 125	
gat gaa tct gat gaa ctg gtc act gat ttt ccc acg gac ctg cca gca	493
Asp Glu Ser Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala	
130 135 140	
acc gaa gtt ttc act cca gtt gtc ccc aca gta gac aca tat gat ggc	541
Thr Glu Val Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly	
145 150 155	
cga ggt gat agt gtg gtt tat gga ctg agg tca aaa tct aag aag ttt	589
Arg Gly Asp Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe	
160 165 170	
cgc aga cct gac atc cag tac cct gat gct aca gac gag gac atc acc	637
Arg Arg Pro Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr	
175 180 185 190	
tca cac atg gaa agc gag gag ttg aat ggt gca tac aag gcc atc ccc	685
Ser His Met Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro	
195 200 205	
gtt gcc cag gac ctg aac gcg cct tct gat tgg gac agc cgt ggg aag	733
Val Ala Gln Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys	
210 215 220	
gac agt tat gaa acg agt cag ctg gat gac cag agt gct gaa acc cac	781
Asp Ser Tyr Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His	
225 230 235	
agc cac aag cag tcc aga tta tat aag cgg aaa gcc aat gat gag agc	829
Ser His Lys Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser	
240 245 250	
aat gag cat tcc gat gtg att gat agt cag gaa ctt tcc aaa gtc agc	877
Asn Glu His Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser	
255 260 265 270	
cgt gaa ttc cac agc cat gaa ttt cac agc cat gaa gat atg ctg gtt	925
Arg Glu Phe His Ser His Glu Phe His Ser His Glu Asp Met Leu Val	
275 280 285	
gta gac ccc aaa agt aag gaa gaa gat aaa cac ctg aaa ttt cgt att	973
Val Asp Pro Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile	
290 295 300	
tct cat gaa tta gat agt gca tct tct gag gtc aat taa aaggagaaaa	1022
Ser His Glu Leu Asp Ser Ala Ser Ser Glu Val Asn	
305 310 315	
aatacaatttt ctcaactttgc atttagtcaa aagaaaaaat gctttatagc aaaatgaaag	1082
agaacatgaa atgcttcttt ctcaagtttat tggttgaatg tgtatctatt tgagtctgga	1142

aataactaat gtgtttgata attagtttag tttgtggctt catggaaact ccctgtaaac 1202
 taaaagcttc agggttatgt ctatgttcat tctatagaag aaatgcaaac tatcactgta 1262
 ttttaatatatt tgttattctc tcatgaatag aaatttatgt agaagcaaac aaaatacttt 1322
 taccactta aaaagagaat ataacatttt atgtcactat aatcttttgt tttttaagtt 1382
 agtgtatatatt ttgttgtgat tatctttttg tggtgtgaat aa 1424

<210> 14
 <211> 314
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Arg Ile Ala Val Ile Cys Phe Cys Leu Leu Gly Ile Thr Cys Ala
 1 5 10 15
 Ile Pro Val Lys Gln Ala Asp Ser Gly Ser Ser Glu Glu Lys Gln Leu
 20 25 30
 Tyr Asn Lys Tyr Pro Asp Ala Val Ala Thr Trp Leu Asn Pro Asp Pro
 35 40 45
 Ser Gln Lys Gln Asn Leu Leu Ala Pro Gln Asn Ala Val Ser Ser Glu
 50 55 60
 Glu Thr Asn Asp Phe Lys Gln Glu Thr Leu Pro Ser Lys Ser Asn Glu
 65 70 75 80
 Ser His Asp His Met Asp Asp Met Asp Asp Glu Asp Asp Asp Asp His
 85 90 95
 Val Asp Ser Gln Asp Ser Ile Asp Ser Asn Asp Ser Asp Asp Val Asp
 100 105 110
 Asp Thr Asp Asp Ser His Gln Ser Asp Glu Ser His His Ser Asp Glu
 115 120 125
 Ser Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala Thr Glu
 130 135 140
 Val Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly Arg Gly
 145 150 155 160
 Asp Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe Arg Arg
 165 170 175
 Pro Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr Ser His
 180 185 190
 Met Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro Val Ala
 195 200 205
 Gln Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys Asp Ser
 210 215 220
 Tyr Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His Ser His
 225 230 235 240
 Lys Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser Asn Glu
 245 250 255
 His Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser Arg Glu
 260 265 270
 Phe His Ser His Glu Phe His Ser His Glu Asp Met Leu Val Val Asp
 275 280 285
 Pro Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile Ser His
 290 295 300
 Glu Leu Asp Ser Ala Ser Ser Glu Val Asn
 305 310

<210> 15

```
<220>
<221> CDS
<222> (100)..(1641)
```

tgggcataga tttcctcttt gagaacatca acctgatttt tgagactttt tqaaaaaatt 60

gac tcc act caa gag agt gag agt gaa gag cac tgg gtg gga ggt ggc 690
Asp Ser Thr Gln Glu Ser Glu Ser Glu Glu His Trp Val Gly Gly Gly

	185	190	195	
agt gat ggg gag agc agc cat gga gac ggc tcc gag ttg gac gat gag				738
Ser Asp Gly Glu Ser Ser His Gly Asp Gly Ser Glu Leu Asp Asp Glu				
	200	205	210	
gga atg cag agt gat gac cca gag agc atc agg agt gaa agg gga aac				786
Gly Met Gln Ser Asp Asp Pro Glu Ser Ile Arg Ser Glu Arg Gly Asn				
	215	220	225	
tcc aga atg aac agt gca ggc atg aaa tca aaa gaa tct gga gaa aac				834
Ser Arg Met Asn Ser Ala Gly Met Lys Ser Lys Glu Ser Gly Glu Asn				
	230	235	240	245
agt gag caa gca aac act caa gat tca ggt ggc agc caa ttg ctg gag				882
Ser Glu Gln Ala Asn Thr Gln Asp Ser Gly Gly Ser Gln Leu Leu Glu				
	250	255	260	
cat ccc agt agg aaa att ttt agg aag tct cgc atc tca gag gaa gat				930
His Pro Ser Arg Lys Ile Phe Arg Lys Ser Arg Ile Ser Glu Glu Asp				
	265	270	275	
gac aga agc gag ctt gat gac aac aac aca atg gaa gaa gtc aag agt				978
Asp Arg Ser Glu Leu Asp Asp Asn Asn Thr Met Glu Glu Val Lys Ser				
	280	285	290	
gac tct aca gaa aac agc aac tcc aga gac act ggc ctc agc caa ccc				1026
Asp Ser Thr Glu Asn Ser Asn Ser Arg Asp Thr Gly Leu Ser Gln Pro				
	295	300	305	
agg aga gac agc aag ggt gac tct caa gaa gac agc aag gag aat ctg				1074
Arg Arg Asp Ser Lys Gly Asp Ser Gln Glu Asp Ser Lys Glu Asn Leu				
	310	315	320	325
tcc cag gaa gag agc caa aac gta gat ggt ccc agc agt gag tcc agc				1122
Ser Gln Glu Glu Ser Gln Asn Val Asp Gly Pro Ser Ser Glu Ser Ser				
	330	335	340	
caa gag gcc aac ctg tca tct caa gag aac agc agt gag tct cag gaa				1170
Gln Glu Ala Asn Leu Ser Ser Gln Glu Asn Ser Ser Glu Ser Gln Glu				
	345	350	355	
gag gtg gtg agt gag tcc agg gga gat aac ccc gac ccc aca act agt				1218
Glu Val Val Ser Glu Ser Arg Gly Asp Asn Pro Asp Pro Thr Thr Ser				
	360	365	370	
tat gta gaa gac cag gaa gac agt gac tcc agc gag gag gac agc tcg				1266
Tyr Val Glu Asp Gln Glu Asp Ser Asp Ser Ser Glu Glu Asp Ser Ser				
	375	380	385	
cac aca ctc tcc cac tca aaa agt gaa tcc aga gag gag caa gca gac				1314
His Thr Leu Ser His Ser Lys Ser Glu Ser Arg Glu Glu Gln Ala Asp				
	390	395	400	405
agc gaa tcc agt gag agc ctc aac ttc tca gag gaa agc ccg gag tcc				1362
Ser Glu Ser Ser Glu Ser Leu Asn Phe Ser Glu Glu Ser Pro Glu Ser				
	410	415	420	
cct gag gat gag aac agc tcc agc cag gag ggc ctc cag tct cac agc				1410
Pro Glu Asp Glu Asn Ser Ser Ser Gln Glu Gly Leu Gln Ser His Ser				
	425	430	435	

```

agc tca gca gag agt cag agc gag gaa agc cat tct gag gaa gac gac 1458
Ser Ser Ala Glu Ser Gln Ser Glu Glu Ser His Ser Glu Glu Asp Asp
      440                      445                      450

agt gac tct caa gac agc agc aga tcc aaa gaa gat agc aac tcc acg 1506
Ser Asp Ser Gln Asp Ser Ser Arg Ser Lys Glu Asp Ser Asn Ser Thr
      455                      460                      465

gag agc aaa tca agc agt gag gaa gat ggc cag ttg aaa aac att gag 1554
Glu Ser Lys Ser Ser Ser Glu Glu Asp Gly Gln Leu Lys Asn Ile Glu
470                      475                      480                      485

ata gag agc cgg aaa tta aca gtt gat gcc tat cac aac aaa ccc att 1602
Ile Glu Ser Arg Lys Leu Thr Val Asp Ala Tyr His Asn Lys Pro Ile
      490                      495                      500

ggg gac caa gat gac aat gac tgc caa gac ggc tat tag catcagctgt 1651
Gly Asp Gln Asp Asp Asn Asp Cys Gln Asp Gly Tyr
      505                      510

cctaagaagc agttgtcaca taaaggagtc ttagggactt gaaaatgtat catgataact 1711

ataatttatt gatgttttga tcaaaagaat aaccagatgc catatttttc ctgaaaggaa 1771

ttgctggaca ttacacttgt ttttaggggtg tcatcatttc acagagggtt aaatactgtg 1831

gagtgcaccc agtaacacag ccaaagagggc tagtaagcaa gaaaggatct gcatgataac 1891

tttgcagctg agatagttcc taattcatca acgtaacaaa caaagctatt ggggtgtccat 1951

gtatatacca ggcactatgc taggtgttga gaatgtaaag cagggttaaga cttgtttctt 2011

gctcctgagg agcgtataga ggggaccac gtagctaaca gagttaacag aggggtatgt 2071

ataagagaaa cgtgaatagt tacaaccacg tatgctgagt gccaaggcca gtgaggctat 2131

gaacacgata tcaatctggg gacactgaga aggataacca atgcgttgga gaacgaggga 2191

ggacttcata gcagaagaat tggagttgga aggttgcata gggtttcctc attggagacc 2251

ggggaattct attcttggga taaataatag taaattctat aggattcaat acttagtttg 2311

caaaagcact ttgaaattat ggcacagtga gttgttttgg ataagaaaat ctttttctcc 2371

caattaattt accaattcat cttttttttt ttttgtagaa ctcccataaa catcacctta 2431

ctaactactg gtgatgataa gaaggattgg gtcaggaaga gtgggagaaa gaaattcctc 2491

tttacgtaga tacttttttag ctttattttt tctaaaatca gtttgcgtgc aatgctagaa 2551

aaaaactgtt ctctgagtcc ttacagagc aaaattctgt atgtaagatt caattgattt 2611

ttgacaaata ccatttgaaa tattacctca acataaaata cttgttttgt aataaagatt 2671

ataataccca g 2682

```

<210> 16
 <211> 513
 <212> PRT

<213> Homo sapiens

<400> 16

Met	Lys	Ile	Ser	Ile	Leu	Leu	Met	Phe	Leu	Trp	Gly	Leu	Ser	Cys	Ala
1				5					10					15	
Leu	Pro	Val	Thr	Arg	Tyr	Gln	Asn	Asn	Glu	Ser	Glu	Asp	Ser	Glu	Glu
			20					25					30		
Trp	Lys	Gly	His	Leu	Ala	Gln	Ala	Pro	Thr	Pro	Pro	Leu	Glu	Ser	Ser
		35					40					45			
Glu	Ser	Ser	Glu	Gly	Ser	Lys	Val	Ser	Ser	Glu	Glu	Gln	Ala	Asn	Glu
	50					55					60				
Asp	Pro	Ser	Asp	Ser	Thr	Gln	Ser	Glu	Glu	Gly	Leu	Gly	Ser	Asp	Asp
65					70					75					80
His	Gln	Tyr	Ile	Tyr	Arg	Leu	Ala	Gly	Gly	Phe	Ser	Arg	Ser	Thr	Gly
			85						90					95	
Lys	Gly	Gly	Asp	Asp	Lys	Asp	Asp	Asp	Glu	Asp	Asp	Ser	Gly	Asp	Asp
			100					105					110		
Thr	Phe	Gly	Asp	Asp	Asp	Ser	Gly	Pro	Gly	Pro	Lys	Asp	Arg	Gln	Glu
		115					120					125			
Gly	Gly	Asn	Ser	Arg	Leu	Gly	Ser	Asp	Glu	Asp	Ser	Asp	Asp	Thr	Ile
	130					135						140			
Gln	Ala	Ser	Glu	Glu	Ser	Ala	Pro	Gln	Gly	Gln	Asp	Ser	Ala	Gln	Asp
145					150					155					160
Thr	Thr	Ser	Glu	Ser	Arg	Glu	Leu	Asp	Asn	Glu	Asp	Arg	Val	Asp	Ser
				165					170					175	
Lys	Pro	Glu	Gly	Gly	Asp	Ser	Thr	Gln	Glu	Ser	Glu	Ser	Glu	Glu	His
			180					185					190		
Trp	Val	Gly	Gly	Gly	Ser	Asp	Gly	Glu	Ser	Ser	His	Gly	Asp	Gly	Ser
		195					200					205			
Glu	Leu	Asp	Asp	Glu	Gly	Met	Gln	Ser	Asp	Asp	Pro	Glu	Ser	Ile	Arg
	210					215					220				
Ser	Glu	Arg	Gly	Asn	Ser	Arg	Met	Asn	Ser	Ala	Gly	Met	Lys	Ser	Lys
225				230						235					240
Glu	Ser	Gly	Glu	Asn	Ser	Glu	Gln	Ala	Asn	Thr	Gln	Asp	Ser	Gly	Gly
				245					250					255	
Ser	Gln	Leu	Leu	Glu	His	Pro	Ser	Arg	Lys	Ile	Phe	Arg	Lys	Ser	Arg
			260					265					270		
Ile	Ser	Glu	Glu	Asp	Asp	Arg	Ser	Glu	Leu	Asp	Asp	Asn	Asn	Thr	Met
		275					280					285			
Glu	Glu	Val	Lys	Ser	Asp	Ser	Thr	Glu	Asn	Ser	Asn	Ser	Arg	Asp	Thr
	290					295					300				
Gly	Leu	Ser	Gln	Pro	Arg	Arg	Asp	Ser	Lys	Gly	Asp	Ser	Gln	Glu	Asp
305					310					315					320
Ser	Lys	Glu	Asn	Leu	Ser	Gln	Glu	Glu	Ser	Gln	Asn	Val	Asp	Gly	Pro
				325					330					335	
Ser	Ser	Glu	Ser	Ser	Gln	Glu	Ala	Asn	Leu	Ser	Ser	Gln	Glu	Asn	Ser
			340					345					350		
Ser	Glu	Ser	Gln	Glu	Glu	Val	Val	Ser	Glu	Ser	Arg	Gly	Asp	Asn	Pro
		355				360						365			
Asp	Pro	Thr	Thr	Ser	Tyr	Val	Glu	Asp	Gln	Glu	Asp	Ser	Asp	Ser	Ser
	370				375						380				
Glu	Glu	Asp	Ser	Ser	His	Thr	Leu	Ser	His	Ser	Lys	Ser	Glu	Ser	Arg
385					390					395					400
Glu	Glu	Gln	Ala	Asp	Ser	Glu	Ser	Ser	Glu	Ser	Leu	Asn	Phe	Ser	Glu
				405					410					415	
Glu	Ser	Pro	Glu	Ser	Pro	Glu	Asp	Glu	Asn	Ser	Ser	Ser	Gln	Glu	Gly
			420					425					430		
Leu	Gln	Ser	His	Ser	Ser	Ser	Ala	Glu	Ser	Gln	Ser	Glu	Glu	Ser	His
		435					440					445			
Ser	Glu	Glu	Asp	Asp	Ser	Asp	Ser	Gln	Asp	Ser	Ser	Arg	Ser	Lys	Glu
450						455						460			

Asp Ser Asn Ser Thr Glu Ser Lys Ser Ser Ser Glu Glu Asp Gly Gln
 465 470 475 480
 Leu Lys Asn Ile Glu Ile Glu Ser Arg Lys Leu Thr Val Asp Ala Tyr
 485 490 495
 His Asn Lys Pro Ile Gly Asp Gln Asp Asp Asn Asp Cys Gln Asp Gly
 500 505 510
 Tyr

<210> 17
 <211> 8201
 <212> DNA
 <213> Homo sapiens

<220>
 <221> exon
 <222> (2387)..(2437)

<220>
 <221> exon
 <222> (3577)..(3660)

<220>
 <221> exon
 <222> (3794)..(4780)

<220>
 <221> exon
 <222> (5257)..(7896)

<400> 17
 attgtcatgc aaaagtccag gacagtgggc cacttttcagt cttcaaagag aaagataaga 60
 aattctggat ttcaaaaatc cttttgaagc cttttaaggt aagatgaaat atccttttta 120
 ctcaagaacca actgattcat ttagaaagaa ctttgaattt caaagatgaa gccagtttga 180
 ttttaagaag cgagtacccc ttaatgatta gattgtatgc ttcctttttg acttgtcata 240
 ttgatagtat gtataaaaaga taacggacga ttacgacctt aggaagagat agattgggaa 300
 gaagaaagac ctctgtactga aaaattggcc aactgaggtg gaaatttgac aattaactat 360
 ctgggcactt tgattagttt tgataaaaaa tgagataact cagatttcaa aaatccacct 420
 tgggctttca aacaaggctt caattaggct ttgcttttta gtattttatt acttactatt 480
 acttattatt tattgtccca catgaaatga aatttagcaa tcactaatga tgccaaatct 540
 aattgctaaa tgaaatgaag ctaaactctca ttccattagt aacaataaat gaaataatct 600
 gatggagctt cacaaattct gaagtctttg ttccatgctg aggtcacctg ggccattttt 660
 attgtagtct tcgaagtcac tcacctgcct tggaaacggt gataaccatc atggaattgt 720
 tcaggagtgg agctgaaaga gagatgtagt ggtcagattt ctgaactgta gctcagaaac 780
 tggacacgta tcactctggc cttggctgca ggtacctttc cagtatgctg aggctcttcc 840
 aaatcacagt gcagacgggc cttctgcaga gctatgtaat gattaggctt gggactgcaa 900
 agtacaggat aactgtggct tagtaaacag ctggccttca acatctgtgc cccagagctc 960
 tgcattgatac ttgtcctggg gtcacctcag cctcacttga atctatggca tttcagaagg 1020
 agctctagct gttcttggct ttctgttgaa cagctataag aatgagcact tttttccctc 1080
 tcagtagctc tggaactgtg tcatctctcc tgtgagaaaa cgccagtaat tctcatgaca 1140
 gttgatattc agtgaagttt tattatattt tcaactaccac cattaaattc aatcaaagcc 1200
 attttatgac atgcagcatt ataacttata catctggtgg gagttcatga aataggagta 1260
 aaactctcct ttctatcatt acttcaagaa atccaacttg caatataaat taattttttt 1320
 actcacacag attataaaat gtctattcca acttatcaga aacatgtttt agaccatttc 1380
 tgaatttgaa ttcaaaaatt gcaaagtgc tgaagcaatg ctagatattg ctgatagtca 1500
 taaatattta tcaacaacat tcagaaaacg tttttttctg tgctttgcat tggaatacaa 1560
 taatcaccaa gacactctcc tgggcctcag gagcttacag gaaatcaggg caacacataa 1620
 gtaactaggc aatttttaaac agtgcaatgc gttaccagtg agacgtgcaa acttccttgg 1680
 tataaaaagg aaagagatac caaataccct ttgaagtggc gtcagagagg gcgtctcaga 1740

gataattcta	ccaaacttca	ggataatcct	gaggtgcagg	tgttggttatt	attccagggtg	1800
gagggataat	aaacctactt	aaattttctca	agcttacaca	gcaagtagca	ggggtaacat	1860
ttgaacccag	gtctctgaat	acaaaccccg	tattctttcc	actagcgtag	gctccctcat	1920
gttagtaatt	tctttctctt	aaagtctggt	atagctcaat	tctatagatt	tggagtaagg	1980
atgacaagtg	ttttaccttt	gaagcacaat	ttcagcagaa	ttagttagta	cttgattaaa	2040
gctattcaga	agagaaatag	atgttttttac	acccaagaat	tgcagaagaa	caaagttaca	2100
gctatgccct	ttgtacctat	tatgggtgttt	tccttcattg	gcacaggcag	aaaaaaatct	2160
aggaagctac	attagtgtctg	agcctgggtga	tgtccccata	accacaccag	gtatgttctg	2220
gaccatcgta	tgtcttctcg	tgttagatac	atgcttcttg	tccaggaaaa	gggcaaatgc	2280
ttacacatca	aaataatata	gtactatgat	tttcccttta	ctttataagt	aattttgtgc	2340
tgttcctttt	ttatacagcc	attgattatt	attattccta	aagaaaatga	agataattac	2400
atatttttgc	atgtgggcag	tagcatgggc	cattccagta	agtatgcctt	tcttagaaaa	2460
cctcttcact	ttgttatctt	ttttaaccta	acattaatac	aaaatgtagt	gtgtgtgtgt	2520
gtgtgtgtgt	gtgtgtgtgt	gtgcatgtac	atgtgtgtat	atatgtgtgt	gtgtatatat	2580
gtttccttaa	tttttttttaa	caggctgagt	ctaaacattt	agatttgcac	taagggtctt	2640
atgtgatata	tgtgaggttt	caacaaaacc	actccaattc	atcgctctcat	tcctctatag	2700
aaactcatat	ctcgtctgaa	ggattattat	tatttaaaac	atttattcag	attaatttac	2760
acttaatgcc	cagaagtcac	ggagactttg	tccatctttg	cttcatactc	tgtgaatttc	2820
attctaatac	gaacaaagtc	tgtgctgttt	aggaagtttc	caagaaagaa	taataagaaa	2880
aagtagattt	tttttcaaca	tataggagac	taatttttca	ctcagagtta	ttatttatgt	2940
gctcactgtg	gaaaattttg	aatatatgac	gaaaaccaat	aaaaaattga	gaaaattcaa	3000
ccattttataa	ttttactagc	cagccatcat	gtttaacatt	ttcatatgct	ttcataatac	3060
caaacatttg	gtattttatgt	agttgaaaat	gttctcaagt	atttcaaagt	tgctcttgca	3120
gagcacagaa	gtatactagc	gtaatacttg	attttgcttc	tgtgcaggct	ctggtcacgc	3180
ctcctgttct	cttaagagtt	ttcatcagga	ttacacttag	agcgggtttg	tgctagtgca	3240
agaggctttt	tgtagagaaa	caccagaggt	ctatcccctc	gtctttctac	aagactcttt	3300
ccttctacag	ttgagataag	tgggctgac	taacacgtcc	ataaaattgg	taataccaca	3360
gtgaaaaata	tccatgtacc	cagtttaaat	tctacacaag	ccctgtaaga	agccacttct	3420
cttttctatc	tgatttagatc	atactttggc	ctttgtgtta	aacctttctt	cttcatggag	3480
ggaagaatat	ttgtgtgtgt	gtgtgtgtgt	gtgcacgctc	acacacatat	tcacaaataa	3540
gaaccttttc	aatagccagt	attttctact	tggcaggttc	ctcaaagcaa	accactggag	3600
agacatgtcg	aaaaatccat	gaattttgcat	ctcctagcaa	gatcaaagt	gtcagtagac	3660
gtataggatg	taatataattt	cattttatatt	cctatttctg	agttgctaca	ttccattaac	3720
ttctccaaga	ttgcaatttg	ctttccttca	agatcattga	cactcataat	tgattgaatt	3780
gtttcttttt	caggatgagt	taaatgccag	tgggaaccatc	aaagaaagt	gtgtcctggg	3840
gcatgaaggt	gatagaggaa	ggcaagagaa	tacccaagat	ggtcacaagg	gagaagggaa	3900
tggctctaag	tgggcagaag	taggagggaa	gagtttttct	acatattcca	cattagcaaa	3960
cgaagagggg	aatattgagg	gctggaatgg	ggacacagga	aaagcagaaa	catatgggtca	4020
tgatggaata	catgggaaag	aagaaaacat	cacagcaa	ggcatccagg	gacaagtaag	4080
catcattgac	aatgctggag	ccacaaacag	aagcaacact	aatggaaata	ctgataagaa	4140
tacccaaaat	ggggatgttg	gcgatgcagg	tcacaatgag	gatgtcgctg	ttgtccaaga	4200
agatggacct	caagtagctg	gaagcaataa	cagtacagac	aatgaggatg	aaataattga	4260
gaattcctgt	agaaacgagg	gtaatacaag	tgaataaca	cctcagatca	acagcaagag	4320
aatgggact	aaggaagctg	aggtaacacc	aggcactgga	gaagatgctg	gcctggataa	4380
ttccgatggg	agtcctagtg	ggaatggagc	agatgaggat	gaagacgagg	gttctggtga	4440
tgatgaagat	gaagaagcag	ggaatggaaa	agacagtagt	aataacagca	agggccagga	4500
gggccaggac	catgggaaag	aagatgatca	tgatagtagc	ataggtcaaa	attcggatag	4560
taaagaatat	tatgaccctg	aaggcaaaaga	agatcccat	aatgaagttg	atggagacaa	4620
gacctccaag	agtgaggaga	attctgctgg	tattccagaa	gacaatggca	gccaaagaat	4680
agaggacacc	cagaagctca	accatagaga	aagcaaacgc	gtagaaaata	gaatcaccaa	4740
agaatcagag	acacatgctg	ttgggaagag	ccaagataag	gttagtttgt	aaagctgatt	4800
tctttcaatg	gcagttttaa	ttcttcccct	ccatctattg	atgctagcac	aaaaataaac	4860
catgacaagc	atccatgtat	ttttgtatcc	atattacttg	actattttaag	gaaatctaga	4920
gtccttacta	gacttcgaga	tagaacaact	ttaaacatct	tacattttctg	ataacttagt	4980
tataattcta	gaaaagtctt	atgtgaaatc	atggatcccc	atgtaattgt	ttacaaaagt	5040
tcctactggg	taggaatgtg	gatgaatttt	taaggaatct	aagcaccagg	atgctttcaa	5100
ttacagaata	aagcacattt	tcacaaataa	ctgtgaagta	ctagaaatgt	aactcctatc	5160
cctatggcaa	cttttcccag	ttattcttcc	tcagatcaat	gcaatttttg	agcaaatatt	5220
cactagttaa	tcattctttc	ctccatcctt	ccatagggaa	tagaaatcaa	gggtcccagc	5280
agtggcaaca	gaaatattac	caaagaagtt	gggaaaggca	acgaaggtaa	agaggataaa	5340
ggacaacatg	gaatgatctt	gggcaaaaggc	aatgtcaaga	cacaaggaga	ggttgtcaac	5400

```

atagaaggac ctggccaaaa atcagaacca ggaaataaag ttggacacag caatacaggt 5460
agtgacagca atagtgatgg atatgacagt tatgattttg atgataagtc catgcaagga 5520
gatgatccca atagcagtga tgaatcta atgctaatc agaaagtgc 5580
aataacagca gtagccgagg agatgcttct tataactctg atgaatcaa agataatggc 5640
aatggcagtg actcaaaagg agcagaagat gatgacagtg atagcacatc agacactaat 5700
aatagtgaca gtaatggcaa tggtaacaat gggaatgatg acaatgacaa atcagacagt 5760
ggcaaaggta aatcagatag cagtgcagtg gatagtagtg atagcagcaa tagcagtgat 5820
agtagtgaca gcagtgacag tgacagcagt gatagcaaca gtagcagtga tagtgacagc 5880
agtgacagtg acagcagtga tagcagtgac agtgatagta gtgatagcag caatagcagt 5940
gacagtagtg acagcagtga tagcagtgac agtagtgata gtagtgacag cagtgcagc 6000
aagtcagaca gcagcaa atc agagagcgac agcagtgata gtgacagtaa gtcagacagc 6060
agtgacagca acagcagtga cagttagtgac aacagtgata gcagcgacag cagcaatagc 6120
agtaacagca gtgatagtag tgacagcagt gatagcagtg acagcagcag tagcagtgac 6180
agcagcagta gcagtgacag cagcaacagc agtgatagta gtgacagtag tgacagcagc 6240
aatagcagtg agagcagtga tagtagtgac agcagtgata gtgacagcag tgatagtagt 6300
gacagcagta atagtaacag cagcgatagt gacagcagca acagcagcga tagcagtgac 6360
agcagtgata gcagtgacag cagcaacagc agtgacagta gcgatagcag tgacagcagc 6420
aacagcagtg acagcagtga tagcagtgac agcagtgata gtagtgacag cagcaacagc 6480
agtgatagca acgacagcag caatagcagt gacagcagtg atagcagcaa cagcagtgat 6540
agcagcaaca gcagtgatag cagtgatagc agtgacagca gtgatagcga cagcagcaat 6600
agcagtgaca gcagtaatag tagtgacagc agcgatagca gcaacagcag tgatagcagc 6660
gacagcagcg atagcagtga cagcagtgat agcgacagca gcaatagaag tgacagtagt 6720
aatagtagtg acagcagcga tagcagtgac agcagcaaca gcagtgacag cagtgatagt 6780
agtgacagca gtgacagcaa cgaaagcagc aatagcagtg acagcagtga tagcagcaac 6840
agcagtgata gtgacagcag tgatagcagc aacagcagtg acagcagtga tagcagcaac 6900
agcagtgata gcagtgaaag cagtaatagt agtgacaaca gcaatagcag tgacagcagc 6960
aacagcagtg acagcagtga tagcagtgac agcagtaata gtagtgacag cagcaatagc 7020
ggtgacagca gcaacagcag tgacagcagt gatagcaata gcagcgacag cagtgcagc 7080
agcaacagca gcgatagcag tgacagcagt gatagcagtg acagcagtga cagcagtgat 7140
agcagcaaca gcagtgatag cagtgcagc agtgacagca gtgatagcag taatagtagt 7200
gacagcagca acagcagtga cagcagcgat agcagtgaca gcagcgatag cagtgcagc 7260
agtgacagca gcaatagcag tgacagcagt gacagcagcg acagcagtga tagcagtgac 7320
agcagtgga gcagcgacag cagtgatagc agtgacagca gtgatagcag cgatagcagt 7380
gacagcagcg acagcagtga cagcagtgac agcagtgaaa gcagcgacag cagcgatagc 7440
agcgacagca gtgacagcag cgacagcagt gacagcagcg atagcagcga cagcagcgac 7500
agcagcgata gcagtgacag cagcaatagc agtgatagca gcgacagcag tgatagcagt 7560
gacagcagcg acagcagcga tagcagcgac agcagtgata gtagtgatag cagtgcagc 7620
agtgacagca gcgacagcag tgacagcagc gacagcagtg acagcagcga cagcagtgac 7680
agcaatgaaa gcagcgacag cagtgcagc agcgatagca gtgacagcag caacagcagt 7740
gacagcagcg acagcagtga tagcagtgac agcacatctg acagcaatga tgagagtgac 7800
agccagagca agtctggtaa cggtaacaac aatggaagtg acagtgacag tgacagtgaa 7860
ggcagtgaca gtaaccactc aaccagtgat gattagaaca aaagaaaaac ccataagatt 7920
ccttttgtga aaagtgtgtt aatgggatag gaaaaaaga tttccaagaa agtaaagaaa 7980
ggggagaaat aaacataaga cgtatgtaaa caaaaacaac tgggggaatc aaatcaaaca 8040
gttggtattca gaaccaagac ctaactcctg cagagacaga ctctgaatgc atgacctttg 8100
gtacatgcct gttaatatct atgttctgaa aatattttgt taaaagtgtg aatctaaaca 8160
taaaagaaca attaaaatat tctttaatac ttcacacaga a 8201

```

<210> 18

<211> 1253

<212> PRT

<213> Homo sapiens

<400> 18

```

Met Lys Ile Ile Thr Tyr Phe Cys Ile Trp Ala Val Ala Trp Ala Ile
  1             5             10             15

```

```

Pro Val Pro Gln Ser Lys Pro Leu Glu Arg His Val Glu Lys Ser Met
      20             25             30

```

Asn Leu His Leu Leu Ala Arg Ser Asn Val Ser Val Gln Asp Glu Leu
 35 40 45
 Asn Ala Ser Gly Thr Ile Lys Glu Ser Gly Val Leu Val His Glu Gly
 50 55 60
 Asp Arg Gly Arg Gln Glu Asn Thr Gln Asp Gly His Lys Gly Glu Gly
 65 70 75 80
 Asn Gly Ser Lys Trp Ala Glu Val Gly Gly Lys Ser Phe Ser Thr Tyr
 85 90 95
 Ser Thr Leu Ala Asn Glu Glu Gly Asn Ile Glu Gly Trp Asn Gly Asp
 100 105 110
 Thr Gly Lys Ala Glu Thr Tyr Gly His Asp Gly Ile His Gly Lys Glu
 115 120 125
 Glu Asn Ile Thr Ala Asn Gly Ile Gln Gly Gln Val Ser Ile Ile Asp
 130 135 140
 Asn Ala Gly Ala Thr Asn Arg Ser Asn Thr Asn Gly Asn Thr Asp Lys
 145 150 155 160
 Asn Thr Gln Asn Gly Asp Val Gly Asp Ala Gly His Asn Glu Asp Val
 165 170 175
 Ala Val Val Gln Glu Asp Gly Pro Gln Val Ala Gly Ser Asn Asn Ser
 180 185 190
 Thr Asp Asn Glu Asp Glu Ile Ile Glu Asn Ser Cys Arg Asn Glu Gly
 195 200 205
 Asn Thr Ser Glu Ile Thr Pro Gln Ile Asn Ser Lys Arg Asn Gly Thr
 210 215 220
 Lys Glu Ala Glu Val Thr Pro Gly Thr Gly Glu Asp Ala Gly Leu Asp
 225 230 235 240
 Asn Ser Asp Gly Ser Pro Ser Gly Asn Gly Ala Asp Glu Asp Glu Asp
 245 250 255
 Glu Gly Ser Gly Asp Asp Glu Asp Glu Glu Ala Gly Asn Gly Lys Asp
 260 265 270
 Ser Ser Asn Asn Ser Lys Gly Gln Glu Gly Gln Asp His Gly Lys Glu
 275 280 285
 Asp Asp His Asp Ser Ser Ile Gly Gln Asn Ser Asp Ser Lys Glu Tyr
 290 295 300
 Tyr Asp Pro Glu Gly Lys Glu Asp Pro His Asn Glu Val Asp Gly Asp
 305 310 315 320
 Lys Thr Ser Lys Ser Glu Glu Asn Ser Ala Gly Ile Pro Glu Asp Asn
 325 330 335
 Gly Ser Gln Arg Ile Glu Asp Thr Gln Lys Leu Asn His Arg Glu Ser
 340 345 350
 Lys Arg Val Glu Asn Arg Ile Thr Lys Glu Ser Glu Thr His Ala Val

355		360		365
Gly Lys Ser Gln Asp Lys Gly Ile Glu Ile Lys Gly Pro Ser Ser Gly				
370		375		380
Asn Arg Asn Ile Thr Lys Glu Val Gly Lys Gly Asn Glu Gly Lys Glu				
385		390		395
Asp Lys Gly Gln His Gly Met Ile Leu Gly Lys Gly Asn Val Lys Thr				
		405		410
				415
Gln Gly Glu Val Val Asn Ile Glu Gly Pro Gly Gln Lys Ser Glu Pro				
		420		425
				430
Gly Asn Lys Val Gly His Ser Asn Thr Gly Ser Asp Ser Asn Ser Asp				
		435		440
				445
Gly Tyr Asp Ser Tyr Asp Phe Asp Asp Lys Ser Met Gln Gly Asp Asp				
		450		455
				460
Pro Asn Ser Ser Asp Glu Ser Asn Gly Asn Asp Asp Ala Asn Ser Glu				
		465		470
				475
Ser Asp Asn Asn Ser Ser Ser Arg Gly Asp Ala Ser Tyr Asn Ser Asp				
		485		490
				495
Glu Ser Lys Asp Asn Gly Asn Gly Ser Asp Ser Lys Gly Ala Glu Asp				
		500		505
				510
Asp Asp Ser Asp Ser Thr Ser Asp Thr Asn Asn Ser Asp Ser Asn Gly				
		515		520
				525
Asn Gly Asn Asn Gly Asn Asp Asp Asn Asp Lys Ser Asp Ser Gly Lys				
		530		535
				540
Gly Lys Ser Asp Ser Ser Asp Ser Asp Ser Ser Asp Ser Ser Asn Ser				
		545		550
				555
Ser Asp Ser Ser Asp Ser Ser Asp Ser Asp Ser Ser Asp Ser Asn Ser				
		565		570
				575
Ser Ser Asp Ser Asp Ser Ser Asp Ser Asp Ser Ser Asp Ser Ser Asp				
		580		585
				590
Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser				
		595		600
				605
Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Lys Ser				
		610		615
				620
Asp Ser Ser Lys Ser Glu Ser Asp Ser Ser Asp Ser Asp Ser Lys Ser				
		625		630
				635
Asp Ser Ser Asp Ser Asn Ser Ser Asp Ser Ser Asp Asn Ser Asp Ser				
		645		650
				655
Ser Asp Ser Ser Asn Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser				
		660		665
				670
Asp Ser Ser Asp Ser Ser Ser Ser Ser Asp Ser Ser Ser Ser Ser Asp				
		675		680
				685

Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser
 690 695 700
 Ser Glu Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Asp Ser Ser Asp
 705 710 715 720
 Ser Ser Asp Ser Ser Asn Ser Asn Ser Ser Asp Ser Asp Ser Ser Asn
 725 730 735
 Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser
 740 745 750
 Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser
 755 760 765
 Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp
 770 775 780
 Ser Asn Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser
 785 790 795 800
 Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser
 805 810 815
 Asp Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser
 820 825 830
 Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser
 835 840 845
 Asp Ser Ser Asp Ser Asp Ser Ser Asn Arg Ser Asp Ser Ser Asn Ser
 850 855 860
 Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser
 865 870 875 880
 Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Glu Ser Ser Asn Ser Ser Asp
 885 890 895
 Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Asp Ser Ser Asp Ser Ser
 900 905 910
 Asn Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Glu
 915 920 925
 Ser Ser Asn Ser Ser Asp Asn Ser Asn Ser Ser Asp Ser Ser Asn Ser
 930 935 940
 Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser
 945 950 955 960
 Asn Ser Gly Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Asn Ser
 965 970 975
 Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser
 980 985 990
 Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp
 995 1000 1005

Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser
 1010 1015 1020

Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser
 1025 1030 1035 1040

Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser Asp Ser Ser Asp
 1045 1050 1055

Ser Ser Asp Ser Ser Asp Ser Ser Gly Ser Ser Asp Ser Ser Asp Ser
 1060 1065 1070

Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser
 1075 1080 1085

Asp Ser Ser Asp Ser Ser Glu Ser Ser Asp Ser Ser Asp Ser Ser Asp
 1090 1095 1100

Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser
 1105 1110 1115 1120

Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn Ser Ser Asp Ser Ser
 1125 1130 1135

Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp
 1140 1145 1150

Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser
 1155 1160 1165

Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Asn
 1170 1175 1180

Glu Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asn
 1185 1190 1195 1200

Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Ser Asp Ser Thr Ser Asp
 1205 1210 1215

Ser Asn Asp Glu Ser Asp Ser Gln Ser Lys Ser Gly Asn Gly Asn Asn
 1220 1225 1230

Asn Gly Ser Asp Ser Asp Ser Asp Ser Glu Gly Ser Asp Ser Asn His
 1235 1240 1245

Ser Thr Ser Asp Asp
 1250

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/09349

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/53 C12N15/90 C12N5/12 C07K16/18 C07K14/51

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BIOSIS, MEDLINE, EPO-Internal, WPI Data, PAJ, STRAND

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BELLAHCENE ET AL: "Expression of bone matrix proteins in human breast cancer: Potential roles in microcalcification formation and in th genesis of bone metastases"</p> <p>BULLETIN DU CANCER, XX, XX, vol. 84, no. 1, January 1997 (1997-01), pages 17-24, XP002107803 figure 1</p> <p>----</p> <p>-/--</p>	1-17, 19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 September 2000

Date of mailing of the international search report

25/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gundlach, B

INTERNATIONAL SEARCH REPORT

Int. l. Application No
PCT/US 00/09349

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PRITCHARD ET AL: "Is the expression of osteopontin and bone sialoprotein greater in breast cancer bone metastases compared to other metastatic sites" EUROPEAN SYMPOSIUM ON CALCIFIED TISSUES, XX, XX, vol. 20, no. 4S, 25 April 1997 (1997-04-25), page 63S XP002107804 abstract	1-17,19
X	US 5 340 934 A (TERMINE JOHN D ET AL) 23 August 1994 (1994-08-23) column 8, line 43 - line 52; figure 5 & DATABASE WPI Derwent Publications Ltd., London, GB; AN 1990-178641 & US 7 432 044 A (FISHER, L.W. ET AL.), 17 April 1990 (1990-04-17) abstract	23
A	PATENT ABSTRACTS OF JAPAN vol. 1997, no. 09, 30 September 1997 (1997-09-30) & JP 09 138213 A (SANGI CO LTD), 27 May 1997 (1997-05-27) abstract	15-24
X	FISHER, L.W. ET AL.: "Antisera and cDNA probes to human and certain animal model bone matrix noncollagenous proteins" ACTA ORTHOPAEDICA SCANDINAVICA, vol. 266, 1995, pages 61-65, XP000937776	15
A	the whole document	1
T	FEDARKO, NEAL S. ET AL: "Factor H binding to bone sialoprotein and osteopontin enables tumor cell evasion of complement-mediated attack" J. BIOL. CHEM., 275(22), 16666-16672, 2 June 2000 (2000-06-02), XP000938681 page 16667, column 2, line 32 - line 35	1-45
-/--		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09349

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	<p>FEDARKO, N. S. (1) ET AL: "Integrins and factor H mediate bone sialoprotein 's and osteopontin's protective properties in complement attack on susceptible cancer cells."</p> <p>JOURNAL OF BONE AND MINERAL RESEARCH, (SEPT., 1999) VOL. 14, NO. SUPPL. 1 PP. S187. MEETING INFO.: TWENTY-FIRST ANNUAL MEETING OF THE AMERICAN SOCIETY FOR BONE AND MINERAL RESEARCH ST. LOUIS, MISSOURI, USA SEPTEMBER 30-OCTOBER 4 1999 AMERICAN SOCIETY,</p> <p>XP000937682</p> <p>abstract</p> <p>-----</p>	1-45